Information For the User Acebrophylline 200 mg tablets

Abrostep 200 SR tablet

Acebrophylline 200 mg Tablet

Composition:

Each uncoated tablet contains:

• **Acebrophylline** – 200 mg

Pharmacological Action:

Acebrophylline is a bronchodilator and mucoregulator with combined properties of ambroxol and theophylline-7-acetate. Its therapeutic actions include:

- Bronchodilation: Relaxes bronchial smooth muscles to relieve bronchospasm and improve airflow in patients with obstructive respiratory conditions.
- Mucoregulation: Lowers mucus viscosity and enhances mucus clearance, helping reduce congestion and making breathing easier.
- Anti-inflammatory: Inhibits the release of pro-inflammatory mediators, reducing inflammation in the airways and respiratory tract.

Indications:

Acebrophylline 200 mg tablets are indicated for the management of:

- Chronic Obstructive Pulmonary Disease (COPD)
- Chronic Bronchitis
- Asthma
- Other Obstructive Airway Diseases

Dosage and Administration:

• **Recommended Dose**: One tablet once or twice daily, or as directed by the physician.

- Administration: Take the tablet with water after meals to minimize gastrointestinal discomfort.
- Duration of Use: Follow the treatment plan as prescribed by your healthcare provider.
 Do not stop the medication abruptly without consulting your doctor.

Precautions:

- Pregnancy & Lactation: Should be used only if prescribed, as the safety of acebrophylline in pregnancy and lactation is not fully established.
- Patients with Liver or Kidney Impairment: Dose adjustments may be necessary; consult with a healthcare provider for guidance.
- Use with Caution: Patients with cardiovascular conditions should use acebrophylline carefully, as it may occasionally lead to palpitations.

Contraindications:

- Known hypersensitivity to acebrophylline, ambroxol, theophylline, or other methylxanthine derivatives.
- Patients with severe hepatic or renal impairment, unless monitored closely by a healthcare provider.

Adverse Reactions:

- Common: Nausea, vomiting, stomach discomfort, headache, and dizziness.
- **Serious but Rare**: Palpitations, skin rashes, difficulty breathing, and other signs of severe allergic reactions. Report any serious side effects to your healthcare provider immediately.

Drug Interactions:

- Avoid combining with other methylxanthines (e.g., theophylline) to reduce the risk of toxicity.
- Caution with other bronchodilators or antihistamines, as they may enhance side effects.

Storage:

- Store in a cool, dry place, below 25°C, protected from light and moisture.
- Keep out of reach of children.

Note: This information is for guidance only. Always follow the directions of a healthcare provider regarding the use of this medication based on individual needs and health status.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for