Information For the Use Amorolfine I.P. 0.25% w/w Cream

Amrofly Cream

Amorolfine I.P. 0.25% w/w Cream

Amorolfine cream is an antifungal medication used to treat a wide range of fungal skin infections. It works by inhibiting the growth of fungi, effectively managing symptoms and promoting healing.

Composition:

• Amorolfine I.P. 0.25% w/w: A topical antifungal agent belonging to the morpholine class, which disrupts fungal cell membrane synthesis.

Uses:

- Fungal Skin Infections:
 - Athlete's foot (Tinea pedis)
 - Ringworm (Tinea corporis)
 - Jock itch (Tinea cruris)
 - Skin candidiasis
- Other Dermatophyte or Yeast Infections:
 - Pityriasis versicolor
 - Nail fold infections caused by fungi

Mechanism of Action:

Amorolfine inhibits ergosterol synthesis, a vital component of fungal cell membranes. This results in weakened membranes, inhibiting fungal growth and leading to their eventual death. **Dosage & Administration:**

- How to Use:
 - 1. Clean and dry the affected area thoroughly before application.
 - 2. Apply a thin layer of the cream to the infected area and surrounding skin.
 - 3. Gently rub it in until fully absorbed.
- Frequency: Use once or twice daily or as directed by your healthcare provider.
- Duration: Continue treatment as prescribed, even if symptoms improve, to prevent recurrence.

Contraindications:

Hypersensitivity to Amorolfine or any of the ingredients in the formulation.

Side Effects:

- Mild irritation, redness, or itching at the site of application.
- Rarely, allergic reactions such as rash or swelling.

Precautions:

- For external use only. Avoid contact with eyes, mouth, or mucous membranes.
- Discontinue use and consult a doctor if severe irritation or allergic reaction occurs.
- Wash hands thoroughly after applying the cream.

• Do not use on open wounds or broken skin.

Storage:

- Store at a temperature below 30°C in a cool, dry place.
- Keep away from direct sunlight and moisture.
- Keep out of reach of children.

Note: Use under medical guidance. If there is no improvement in symptoms after 2-4 weeks of use, consult your healthcare provider.

Manufactured in India for:

TM

Cafoli Lifecare Pvt. Ltd. (An ISO 9001: 2015 Certified Co.) Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113 TM: Trademark Applied for