

# Cefzlan 200 Dry Syrup

**Brand Name:** Cefzlan 200 Dry Syrup

**Dosage Form:** Dry Syrup

**Composition:**

After reconstitution, each 5 ml contains:

- Cefixime IP 200 mg
  - Flavoured syrup base q.s.
- 

**Description:**

Cefzlan 200 Dry Syrup contains Cefixime, a third-generation cephalosporin antibiotic with broad-spectrum antibacterial activity. It is indicated for the treatment of various bacterial infections caused by susceptible organisms. It helps in inhibiting bacterial cell wall synthesis, thereby controlling infection and promoting recovery.

---

**Indications:**

Indicated for the treatment of:

- Respiratory tract infections
- Urinary tract infections
- Typhoid fever
- Otitis media
- Pharyngitis and tonsillitis

It helps in reducing bacterial load, relieving symptoms, and preventing progression of infection.

---

**Dosage & Administration:**

Dosage should be as directed by the physician based on age, body weight, and severity of infection.

Reconstitute the dry syrup with the recommended quantity of boiled and cooled water up to the marked level and shake well before use.

Shake the bottle thoroughly before each administration.

Complete the full prescribed course of therapy.

---

**Pharmacological Action:**

- Cefixime is a bactericidal cephalosporin antibiotic that inhibits bacterial cell wall synthesis by binding to penicillin-binding proteins (PBPs).
- This results in weakening of the bacterial cell wall and subsequent bacterial cell death.

These actions help in effective eradication of susceptible bacterial pathogens.

---

**Side Effects:**

Common side effects include diarrhea, nausea, abdominal pain, indigestion, and headache. Rarely, allergic reactions, skin rash, or changes in liver enzymes may occur.

---

**Precautions & Warnings:**

- Use with caution in patients with penicillin or cephalosporin hypersensitivity

- **Dose adjustment may be required in renal impairment**
  - **Prolonged use may lead to superinfection or resistant organisms**
  - **Discontinue use if severe allergic reaction occurs**
- 

**Drug Interactions:**

**May interact with anticoagulants and nephrotoxic drugs.**

**Concomitant use with certain medications may alter renal elimination.**

---

**Storage:**

**Store the dry syrup below 25°C in a cool, dry place.**

**After reconstitution, store as directed and use within the prescribed period.**

**Protect from light and keep out of reach of children.**

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

**Corporate Office :**

Plot no. 307, Industrial Area Phase-2,  
Panchkula, Haryana -134113

TM: Trademark Applied for