

Information For the User
Cefixime (200mg) + Dicloxacillin (500mg) ER Tablet

Cefzlan DC Tablet

Brand Name: Cefzlan DC Tablet

Composition:

Each extended-release tablet contains:

Cefixime IP 200 mg

Dicloxacillin Sodium IP equivalent to Dicloxacillin 500 mg (Extended Release)

Therapeutic Indication:

Indicated in the treatment of respiratory tract infections, skin and soft tissue infections, ENT infections, urinary tract infections, and dental infections

Effective against mixed gram-positive and gram-negative bacterial infections, particularly where β -lactamase-producing organisms are suspected

Mechanism of Action:

Cefixime is a third-generation cephalosporin antibiotic that inhibits bacterial cell wall synthesis by binding to penicillin-binding proteins

Effective mainly against gram-negative organisms

Dicloxacillin is a penicillinase-resistant penicillin that is effective against gram-positive cocci, including penicillinase-producing *Staphylococcus aureus*

The combination offers broad-spectrum antibacterial activity with β -lactamase resistance

Dosage & Administration:

Usual adult dose: One tablet once or twice daily or as directed by the physician

Swallow whole with water, preferably after meals

Duration of therapy depends on severity and type of infection

Contraindications:

Hypersensitivity to cephalosporins, penicillins, or any component of the formulation

History of severe allergic reactions (e.g., anaphylaxis) to β -lactam antibiotics

Neonates or patients with hepatic dysfunction (for Dicloxacillin use)

Warnings & Precautions:

Use with caution in patients with renal or hepatic impairment

Assess for any cross-sensitivity in patients with penicillin allergy

Prolonged use may result in superinfection with resistant organisms including fungi

Monitor for signs of gastrointestinal disturbances or pseudomembranous colitis

Side Effects:

Common: nausea, vomiting, diarrhea, abdominal pain

Less common: rash, urticaria, headache, dizziness

Rare: anaphylaxis, Stevens-Johnson syndrome, hepatic dysfunction

Storage:

Store in a cool, dry, and dark place below 25°C

Protect from moisture and direct sunlight
Keep out of reach of children

Packaging:

Cefzlan DC Tablet is available in alu-alu or blister strip packs

Schedule:

Schedule H Prescription Drug – To be sold by retail on the prescription of a registered medical practitioner only

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for