

Chymoedge AP Tablet

Aceclofenac (100 mg) + Paracetamol (325 mg) + Trypsin Chymotrypsin (50000 AU) Tablet

reduce swelling and inflammation in tissues.

Composition

Each tablet contains:

- **Aceclofenac: 100 mg**
 - **Paracetamol (Acetaminophen): 325 mg**
 - **Trypsin Chymotrypsin: 50000 AU (activity unit)**
-

Description

This combination tablet contains Aceclofenac, a non-steroidal anti-inflammatory drug (NSAID), Paracetamol, an analgesic and antipyretic, and Trypsin Chymotrypsin, a proteolytic enzyme combination used for its anti-inflammatory properties. The combination provides effective relief from pain and inflammation, especially in conditions like musculoskeletal disorders, acute injuries, and postoperative pain.

- **Aceclofenac works by inhibiting the synthesis of prostaglandins, which are chemicals responsible for inflammation, pain, and fever.**
- **Paracetamol helps reduce fever and relieve mild to moderate pain, such as headaches, muscle pain, or toothaches.**
- **Trypsin Chymotrypsin has an enzyme activity that assists in the breakdown of proteins and can help**

Indications

This combination is indicated for the relief of:

- **Acute pain (such as musculoskeletal pain, dental pain, or post-surgical pain)**
 - **Chronic pain (in conditions like osteoarthritis, rheumatoid arthritis)**
 - **Inflammatory conditions (like soft tissue injuries or sprains)**
 - **Fever associated with pain**
-

Dosage and Administration

- **Adults: Take 1 tablet twice daily after meals.**
- **Children: The use of this combination in children is not recommended unless prescribed by a healthcare provider.**

Administration:

- **Swallow the tablet whole with a glass of water.**
 - **It is advised to take the tablet after meals to minimize stomach irritation, especially due to Aceclofenac.**
-

Mechanism of Action

- **Aceclofenac is an NSAID that inhibits cyclooxygenase (COX-1 and COX-2), enzymes responsible for producing prostaglandins, leading to reduced pain, swelling, and inflammation.**
 - **Paracetamol works centrally in the brain to reduce the perception of pain and regulate body temperature, offering relief from mild to moderate pain and fever.**
 - **Trypsin Chymotrypsin breaks down proteins involved in the inflammatory process, thus reducing swelling and aiding tissue repair.**
-

Warnings and Precautions

- **Gastrointestinal: Long-term use of Aceclofenac may cause gastric irritation or ulcers. It is advisable to use this combination for the shortest duration possible.**
 - **Liver or kidney disease: Caution is advised if you have pre-existing liver or kidney conditions, as both Aceclofenac and Paracetamol are metabolized in the liver.**
 - **Cardiovascular disease: NSAIDs like Aceclofenac should be used cautiously in individuals with cardiovascular disease, as they can increase the risk of heart attack or stroke.**
 - **Alcohol: Avoid or limit alcohol consumption as it can increase the risk of liver damage when taken with Paracetamol and may exacerbate gastrointestinal side effects with Aceclofenac.**
-

Side Effects

Common side effects may include:

- **Gastrointestinal: Nausea, dyspepsia (indigestion), abdominal pain, diarrhea, or constipation.**
 - **Headache**
 - **Dizziness or drowsiness**
 - **Rash**
 - **Swelling (due to Trypsin Chymotrypsin's action on protein breakdown)**
-

Storage

- **Store in a cool, dry place below 25°C.**
 - **Keep the tablets in their original packaging to protect from moisture.**
 - **Keep out of reach of children.**
-

Note: Always follow your doctor's instructions on how to use this medication. Do not exceed the recommended dose. If you experience any side effects, or if symptoms persist, consult your healthcare provider.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for