

Information For the Use of  
Ciprofloxacin (0.3% w/v) Eye Drops

# Ciproplit Eye Drops

**Brand Name:** Ciproplit Eye Drops

**Dosage Form:** Eye Drops

**Composition:**

**Each ml contains:**

- Ciprofloxacin Hydrochloride IP equivalent to Ciprofloxacin 0.3% w/v
- Sterile ophthalmic vehicle q.s.

---

**Description:**

Ciproplit Eye Drops contains Ciprofloxacin, a broad-spectrum fluoroquinolone antibacterial agent used for the treatment of bacterial infections of the eye caused by susceptible microorganisms. It helps in eliminating pathogenic bacteria, reducing infection-related symptoms, and promoting healing of ocular tissues.

---

**Indications:**

Indicated for the treatment of:

- Bacterial conjunctivitis
- Corneal ulcers caused by susceptible bacteria
- Blepharoconjunctivitis
- Other ophthalmic bacterial infections as determined by the physician

It helps in controlling bacterial growth, relieving ocular discomfort, and preventing the spread of infection.

---

**Dosage & Administration:**

Dosage should be as directed by the physician.

Instill the prescribed number of drops into the affected eye(s).

Wash hands before administration and avoid touching the dropper tip to any surface.

Complete the full course of treatment as prescribed.

---

**Pharmacological Action:**

- Ciprofloxacin inhibits bacterial DNA gyrase and topoisomerase IV, enzymes essential for bacterial DNA replication, transcription, and repair.
- This results in inhibition of bacterial growth and eventual bacterial cell death.

These actions provide effective treatment against a wide range of susceptible ocular bacterial pathogens.

---

**Side Effects:**

Common side effects include transient burning or stinging sensation, eye discomfort, redness, itching, blurred vision, and foreign body sensation.

Rarely, hypersensitivity reactions or white crystalline deposits on the cornea may occur during treatment of corneal ulcers.

---

**Precautions & Warnings:**

- For ophthalmic use only
- Not effective against viral or fungal eye infections
- Discontinue use if signs of hypersensitivity occur
- Avoid wearing contact lenses during active eye infection unless advised by the physician
- Avoid contamination of the dropper tip to prevent secondary infection

---

**Drug Interactions:**

No significant systemic interactions are expected due to minimal systemic absorption.

If used with other ophthalmic medications, maintain an interval of at least 5–10 minutes between applications.

---

**Storage:**

**Store below 25°C in a cool, dry place.**  
**Protect from light.**  
**Do not freeze.**  
**Keep the bottle tightly closed after use.**  
**Keep out of reach of children.**

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

**Corporate Office :**

Plot no. 307, Industrial Area Phase-2,  
Panchkula, Haryana -134113

TM: Trademark Applied for