

Information For the Use of
Ciprofloxacin (0.3% w/v) + Dexamethasone (0.1% w/v) + Benzalkonium Chloride Solution IP 0.02% w/v
Eye Drops

Ciproplit D Eye Drops

Brand Name: Ciproplit D Eye Drops

Dosage Form: Eye Drops

Composition:

Each ml contains:

- Ciprofloxacin Hydrochloride IP equivalent to Ciprofloxacin 0.3% w/v
- Dexamethasone Sodium Phosphate IP equivalent to Dexamethasone 0.1% w/v
- Benzalkonium Chloride Solution IP 0.02% w/v (as preservative)
- Sterile ophthalmic vehicle q.s.

Description:

Ciproplit D Eye Drops is a sterile ophthalmic preparation containing Ciprofloxacin, a broad-spectrum fluoroquinolone antibiotic, and Dexamethasone, a potent corticosteroid. It is indicated for steroid-responsive ocular inflammatory conditions where bacterial infection or the risk of bacterial infection exists. It helps in controlling bacterial growth while simultaneously reducing ocular inflammation, redness, swelling, and discomfort.

Indications:

Indicated for the treatment of:

- Bacterial eye infections associated with inflammation
- Bacterial conjunctivitis with inflammatory symptoms
- Post-operative ocular inflammation with risk of bacterial infection
- Steroid-responsive ocular inflammatory conditions requiring antibacterial coverage

It helps in eliminating susceptible bacteria and reducing inflammation-related ocular symptoms.

Dosage & Administration:

Dosage should be as directed by the physician.

Instill the prescribed number of drops into the affected eye(s).

Wash hands thoroughly before administration and avoid touching the dropper tip to any surface.

If using multiple ophthalmic medications, maintain an interval of at least 5–10 minutes between applications.

Pharmacological Action:

- Ciprofloxacin inhibits bacterial DNA gyrase and topoisomerase IV, preventing bacterial replication and leading to bacterial cell death.
- Dexamethasone suppresses inflammatory responses by inhibiting inflammatory mediators, reducing edema, redness, and tissue irritation.

Together, these actions provide effective antibacterial and anti-inflammatory therapy for ocular conditions.

Side Effects:

Common side effects include transient burning, stinging, eye irritation, redness, blurred vision, and discomfort after instillation.

Prolonged corticosteroid use may increase intraocular pressure, delay wound healing, or increase susceptibility to secondary infections.

Rarely, hypersensitivity reactions may occur.

Precautions & Warnings:

- For ophthalmic use only
- Not indicated for viral or fungal eye infections
- Prolonged use may result in increased intraocular pressure or glaucoma
- Monitor intraocular pressure during extended treatment

- Remove contact lenses before administration and reinsert only as advised by the physician
- Avoid contamination of the dropper tip

Drug Interactions:

No significant systemic drug interactions are expected due to minimal systemic absorption.

Concurrent use with other ophthalmic corticosteroids may increase the risk of ocular adverse effects.

Storage:

Store below 25°C in a cool, dry place.

Protect from light.

Do not freeze.

Keep the bottle tightly closed after use.

Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113

TM: Trademark Applied for