

# Clotimet CTP Tablet

## Clotrimazole, Tinidazole & Povidone Iodine Vaginal Tablet

to reduce infection risk by killing bacteria, viruses, and fungi.

---

### Formulation:

Each tablet contains:

- **Clotrimazole**
- **Tinidazole**
- **Povidone Iodine**

### Strength:

- Clotrimazole: 100 mg
- Tinidazole: 500 mg
- Povidone Iodine: 100 mg

### Type:

Vaginal tablet

---

### Uses:

This combination tablet is used for the treatment of various vaginal infections, including:

1. **Fungal Infections (Vaginal Candidiasis):**
  - Clotrimazole, an **antifungal** agent, is effective against yeast infections (candidiasis), commonly caused by *Candida* species.
2. **Bacterial Infections:**
  - Tinidazole, an **antiprotozoal** and **antibacterial** agent, treats infections caused by anaerobic bacteria and protozoa, such as **bacterial vaginosis** and **trichomoniasis**.
3. **Antiseptic Action:**
  - Povidone Iodine provides broad-spectrum **antiseptic** properties, helping

---

### Mechanism of Action:

- **Clotrimazole** inhibits the growth of fungi by damaging their cell membranes, which prevents their ability to reproduce and thrive.
- **Tinidazole** interferes with the DNA of susceptible microorganisms, leading to their death and preventing further infection.
- **Povidone Iodine** releases iodine in the affected area, exerting a broad antimicrobial action against bacteria, viruses, and fungi.

---

### Dosage:

- **Typical Dose:**
  - Insert **one vaginal tablet** once daily, preferably at bedtime, for a duration of **3-7 days** as prescribed by a healthcare provider.
- **Administration:**
  - The tablet should be inserted **deeply into the vagina**, preferably using a clean applicator if provided.
  - If you are unsure of how to insert the tablet, consult your healthcare provider.

---

### Precautions:

- **Pregnancy and Breastfeeding:**
  - Consult a doctor before use if you are pregnant, planning to become pregnant, or breastfeeding.
- **Allergic Reactions:**
  - If you experience any signs of an allergic reaction, such as rash, itching, swelling,

or difficulty breathing, seek medical attention immediately.

- **Pre-existing Conditions:**
  - Inform your doctor if you have any history of **liver disease, kidney disease, or vaginal ulcers.**
- **Avoid Douching:**
  - Do not douche the vagina during treatment, as it can interfere with the effectiveness of the medication.

- Store at **20°C to 25°C** (68°F to 77°F).
- Keep the tablets in their original packaging, protected from moisture and light.
- Keep out of reach of children.

---

## Side Effects:

- **Common:**
  - Vaginal irritation
  - Itching or burning sensation
  - Increased vaginal discharge
  - Abdominal discomfort
- **Less Common:**
  - Nausea
  - Headache
  - Skin rash
- **Serious (Seek Medical Attention):**
  - Severe allergic reactions (swelling of the face, lips, or throat, difficulty breathing)
  - Severe vaginal irritation or pain

---

## Note:

- This medication is for **vaginal use only** and should not be taken orally.
- Complete the full course of treatment, even if symptoms improve before the treatment period ends. Discontinuing early may result in the recurrence of infection.
- If symptoms persist or worsen after completing the treatment, consult your healthcare provider for further evaluation.

Manufactured in India for:

**Cafoli**<sup>TM</sup>  
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for

---

## Drug Interactions:

- **Alcohol:**
  - Avoid alcohol consumption while using **Tinidazole**, as it can cause a **disulfiram-like reaction**, leading to symptoms like nausea, vomiting, and flushing.
- **Other Vaginal Medications:**
  - Inform your doctor if you are using other vaginal products to avoid potential interactions.
- **Other Drugs:**
  - Always inform your healthcare provider about any other medications, supplements, or herbs you are taking to avoid interactions.

---

## Storage: