

Information For the User
Beclometasone (0.025% w/v) + Clotrimazole (1% w/v) lotion

Clotimet B Lotion

Product Name: Clotimet B Lotion

Composition: Beclometasone Dipropionate I.P. 0.025% w/v + Clotrimazole I.P. 1% w/v

Dosage Form: Topical Lotion

Description:

Clotimet B Lotion is a topical antifungal and anti-inflammatory preparation containing Clotrimazole and Beclometasone Dipropionate. Clotrimazole is an imidazole antifungal agent that interferes with fungal cell membrane permeability, while Beclometasone is a potent corticosteroid that reduces inflammation, redness, and itching.

Composition (per ml):

- **Clotrimazole I.P.** – 1% w/v
- **Beclometasone Dipropionate I.P.** – 0.025% w/v
- **Excipients** – q.s.

Indications:

Clotimet B Lotion is indicated for the topical treatment of:

1. Fungal infections of the skin such as tinea corporis, tinea cruris, and tinea pedis
2. Inflammatory dermatoses associated with secondary fungal infections
3. Eczematous dermatitis with fungal superinfection

Mechanism of Action:

- **Clotrimazole:** Acts by inhibiting ergosterol synthesis, an essential component of the fungal cell membrane, causing increased permeability and cell death.
- **Beclometasone:** Binds to glucocorticoid receptors, reducing the formation of inflammatory mediators and alleviating symptoms like itching and redness.

Dosage and Administration:

- Clean and dry the affected area.
- Apply a thin layer of lotion to the affected area twice daily (morning and evening), or as directed by a physician.
- Do not use for more than 2 weeks continuously without medical supervision.

Side Effects:

- **Common:** Burning sensation, itching, irritation, or redness at the application site
 - **Rare:** Skin thinning, striae, or contact dermatitis with prolonged use
-

Precautions:

1. **External Use Only:** Not to be used in eyes or ingested.
2. **Hypersensitivity:** Discontinue use if signs of allergic reaction occur.
3. **Pediatric Use:** Use with caution under medical supervision.
4. **Prolonged Use:** Avoid continuous use over large areas to prevent systemic absorption of corticosteroids.

Contraindications:

- Known hypersensitivity to clotrimazole, beclometasone, or any component of the formulation
- Untreated bacterial, viral, or tubercular skin infections

Storage:

- Store below 25°C in a cool, dry place
- Protect from light and moisture
- Keep out of reach of children

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for