Information For the Use Levosalbutamol 100mcg + Ipratropium 40mcg

Critsal Capsule

Levosalbutamol 100mcg + Ipratropium 40mcg Flipcaps: Information Guide the airways. This results in further improvement in airflow and breathing.

Description

Levosalbutamol 100mcg and Ipratropium 40mcg Flipcaps is a combination inhaler designed to manage respiratory conditions such as chronic obstructive pulmonary disease (COPD) and asthma. Levosalbutamol is a selective beta-2 agonist that works as a bronchodilator, while Ipratropium is an anticholinergic agent that helps to open up the airways by inhibiting bronchoconstriction.

Composition (per dose)

- Levosalbutamol: 100mcg (bronchodilator)
- Ipratropium: 40mcg (anticholinergic)

Mechanism of Action

- 1. Levosalbutamol:
 - A beta-2 adrenergic agonist that stimulates beta-2 receptors in the smooth muscle of the airways, leading to muscle relaxation and bronchodilation. This allows easier airflow into the lungs and helps relieve symptoms of bronchospasm.
- 2. Ipratropium:
 - An anticholinergic agent that blocks the action of acetylcholine on muscarinic receptors, preventing bronchoconstriction and reducing mucus production in

Indications

Levosalbutamol 100mcg + Ipratropium 40mcg Flipcaps is indicated for:

- Chronic Obstructive Pulmonary Disease (COPD): To relieve bronchospasm associated with COPD.
- Asthma: As a maintenance treatment for asthma exacerbations and to improve lung function.
- Acute exacerbations of COPD or asthma, when fast relief of bronchoconstriction is needed.

Dosage and Administration

- Adults:
 - Inhale 1 flipcap (100mcg Levosalbutamol + 40mcg Ipratropium) two to four times per day, or as directed by a healthcare provider.
 - Do not exceed the prescribed dose.
 - Children (6 years and older):
 - Inhalation of 1 flipcap one to two times per day, or as prescribed.
 - Consult a healthcare provider for proper dose adjustment for children.

How to Use:

• Insert the flipcap into the inhaler device and inhale the medication as directed.

• Ensure you rinse your mouth after use to prevent irritation and dry mouth.

Contraindications

- Hypersensitivity to Levosalbutamol, Ipratropium, or any other components of the flipcap.
- Severe hypersensitivity to atropine or its derivatives.
- Patients with a history of narrow-angle glaucoma, urinary retention, or prostate enlargement should use this medication with caution.

Warnings and Precautions

- Caution in Cardiovascular Disease: Levosalbutamol may increase heart rate and blood pressure. Use with caution in patients with a history of heart disease or hypertension.
- Glaucoma: Ipratropium can cause increased intraocular pressure. Avoid use in patients with glaucoma or a predisposition to glaucoma.
- Urinary Retention: Use with caution in patients with bladder neck obstruction or prostate hypertrophy.
- Paradoxical Bronchospasm: Rarely, inhalation of beta-agonists may cause paradoxical bronchospasm, which can be life-threatening. Discontinue use if this occurs and seek immediate medical attention.

Adverse Effects

• Common:

- Dry mouth, cough, throat irritation, headache, dizziness, or nausea.
- Less Common:
 - Palpitations, tremors, nervousness, or urinary retention.
- Serious (Rare):
 - Paradoxical bronchospasm, allergic reactions (swelling of the face, lips, or tongue), or blurred vision.
 - Severe tachycardia or arrhythmia.

Drug Interactions

- Beta-blockers: May reduce the effectiveness of Levosalbutamol and increase the risk of bronchospasm.
- Anticholinergic Drugs: Combining with other anticholinergic medications may increase the risk of urinary retention and dry mouth.
- Diuretics: May enhance the risk of hypokalemia when used with bronchodilators.
- Other Respiratory Medications: Use caution when combining with other inhaled bronchodilators or steroids.

Storage

- Store in a cool, dry place, at a temperature not exceeding 25°C.
- Keep away from heat, moisture, and direct sunlight.
- Keep out of reach of children.

Key Points

This combination inhaler provides effective relief for bronchospasm and helps manage symptoms of asthma and COPD.

- Follow the prescribed dosage to avoid overuse or underuse of the medication.
- Proper inhaler technique is important for the effectiveness of the medication.
- If symptoms persist or worsen, consult your doctor immediately.

Note: This medication should be used under the supervision of a healthcare provider. If you experience any severe or unexpected side effects, contact your healthcare provider immediately.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd. (An ISO 9001: 2015 Certified Co.) Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113 TM: Trademark Applied for