

# Cufsniff LD Syrup

## Information for the Use

---

**Chlorpheniramine Maleate (2 mg/5 mL) +  
Levodropropizine (30 mg/5 mL) Syrup**

---

### Composition:

Each 5 mL contains:

- Chlorpheniramine Maleate: 2 mg
  - Levodropropizine: 30 mg
- 

### Description:

This syrup combines two active ingredients:

- Chlorpheniramine Maleate, an antihistamine that relieves symptoms of allergies such as sneezing, runny nose, and itching.
  - Levodropropizine, a cough suppressant that reduces the frequency and intensity of dry, non-productive coughs by acting on the peripheral nervous system without affecting the central nervous system.
- 

### Indications:

**Chlorpheniramine Maleate (2 mg/5 mL) +  
Levodropropizine (30 mg/5 mL) Syrup is  
indicated for:**

- Dry, non-productive cough associated with allergies or respiratory infections.
- Relief from cough accompanied by allergic rhinitis.
- Treatment of cough in conditions like bronchitis or post-viral cough where mucus production is minimal.

## Dosage and Administration:

- **Adults and Children Above 12 Years: 5 mL (1 teaspoon) 2–3 times a day.**
- **Children (6–12 Years): 2.5 mL (½ teaspoon) 2–3 times a day.**
- **Children (2–6 Years): As directed by the physician.**

## Method of Administration:

- Shake the bottle well before use.
  - Administer with a measuring cup or spoon for accurate dosing.
  - Can be taken with or without food.
- 

## Contraindications:

- Hypersensitivity to Chlorpheniramine Maleate, Levodropropizine, or any excipients in the formulation.
  - Severe liver or kidney impairment.
  - Patients with productive cough or chronic respiratory conditions like asthma.
  - Children under 2 years of age.
- 

## Precautions:

- **Drowsiness:** Chlorpheniramine Maleate may cause drowsiness; avoid activities requiring alertness (e.g., driving).
- **Liver or Kidney Disorders:** Use with caution in patients with hepatic or renal impairment.
- **Pregnancy and Lactation:**

- **Pregnancy:** Use only if clearly necessary and as prescribed by a healthcare professional.
  - **Lactation:** Not recommended as it may pass into breast milk.
- 

- **Do not exceed the recommended dosage.**
  - **Report any unusual or severe side effects to a healthcare professional immediately.**
  - **Discontinue use and consult a doctor if the cough persists for more than 7 days or is accompanied by fever, rash, or persistent headache.**
- 

### **Possible Side Effects:**

#### **Common side effects:**

- **Drowsiness or sedation**
- **Dry mouth**
- **Nausea or vomiting**
- **Dizziness**

#### **Rare but serious side effects:**

- **Allergic reactions (rash, itching, swelling)**
  - **Difficulty breathing**
  - **Severe drowsiness**
- 

### **Drug Interactions:**

- **CNS Depressants (e.g., alcohol, sedatives, tranquilizers): May enhance the sedative effects of Chlorpheniramine Maleate.**
  - **Antihypertensive Drugs: May cause additive effects leading to hypotension.**
  - **Avoid concurrent use with other cough suppressants unless directed by a healthcare professional.**
- 

### **Storage:**

- **Store at room temperature below 25°C (77°F).**
  - **Protect from light and moisture.**
  - **Keep out of reach of children.**
- 

### **Special Instructions:**

- **Use the syrup exactly as prescribed by your doctor.**

**Note: This medication is for symptomatic relief and does not treat the underlying cause of the cough. Use only under the supervision of a healthcare professional. Always read the label and package insert for additional information.**

Manufactured in India for:

**Cafoli**<sup>TM</sup>  
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for