

Information For the User
Dapagliflozin 10mg + Metformin Hydrochloride I.P. 1000mg Tablet

Dapoflit M Forte

Brand Name: Dapoflit M Forte Tablet

Dosage Form: Tablet

Composition:

Each film-coated tablet contains:

- Dapagliflozin IP 10 mg
- Metformin Hydrochloride IP 1000 mg

Description:

Dapoflit M Forte Tablet is a fixed-dose combination of Dapagliflozin, a sodium-glucose co-transporter 2 (SGLT2) inhibitor, and Metformin Hydrochloride, a biguanide antidiabetic agent. It is indicated for the management of type 2 diabetes mellitus when glycemic control is inadequate with monotherapy. It helps in improving glycemic control by reducing renal glucose reabsorption and decreasing hepatic glucose production.

Indications:

Indicated for the treatment of:

- Type 2 diabetes mellitus
- Patients inadequately controlled on metformin or SGLT2 inhibitor monotherapy

It helps in controlling fasting and post-prandial blood glucose levels.

Dosage & Administration:

Dosage should be as directed by the physician.

The tablet should be swallowed whole with water and preferably taken with meals to minimize gastrointestinal side effects.

Regular monitoring of blood glucose and renal function is recommended.

Pharmacological Action:

Dapagliflozin inhibits SGLT2 in the proximal renal tubules, leading to increased urinary glucose excretion. Metformin decreases hepatic gluconeogenesis, improves insulin sensitivity, and reduces intestinal glucose absorption, resulting in complementary glucose-lowering effects.

Side Effects:

Generally well tolerated.

Possible side effects include gastrointestinal disturbances, increased urination, urinary tract infections, genital infections, and headache.

Precautions & Warnings:

- Use with caution in patients with renal impairment
- Assess renal function prior to initiation and periodically thereafter
- Risk of lactic acidosis with metformin, especially in predisposed patients
- Not recommended in type 1 diabetes mellitus or diabetic ketoacidosis

Drug Interactions:

Concomitant use with insulin or insulin secretagogues may increase the risk of hypoglycemia. Alcohol may potentiate the risk of lactic acidosis.

Storage:

Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for