Decplist Gel

Diclofenac Diethylamine (1.16% w/v) + Linseed Oil (3% w/v) + Methyl Salicylate (10% w/v) + Menthol (5% w/v) Gel: Information Guide

Description

This topical gel combines the antiinflammatory and analgesic properties of Diclofenac Diethylamine, Linseed Oil, Methyl Salicylate, and Menthol to provide effective relief from pain and inflammation in musculoskeletal and joint conditions.

Composition

- Diclofenac Diethylamine: 1.16% w/v (a non-steroidal anti-inflammatory drug NSAID).
- Linseed Oil: 3% w/v (an antiinflammatory agent).
- Methyl Salicylate: 10% w/v (a counterirritant and analgesic).
- Menthol: 5% w/v (provides a cooling effect and pain relief).
- Dosage Form: Topical gel.

Mechanism of Action

- 1. Diclofenac Diethylamine: Reduces pain and inflammation by inhibiting the cyclooxygenase (COX) enzymes responsible for prostaglandin synthesis.
- 2. Linseed Oil: Contains omega-3 fatty acids that reduce inflammation and enhance skin absorption of active ingredients.

- 3. Methyl Salicylate: Acts as a counterirritant, creating a warming sensation that distracts from deeper musculoskeletal pain.
- 4. Menthol: Provides a cooling effect, reducing pain perception and improving local blood flow.

Indications

This gel is indicated for the symptomatic relief of:

- Muscle pain and sprains.
- Joint pain, stiffness, and inflammation associated with arthritis.
- Backache, neck pain, and shoulder pain.
- Sports injuries like strains and contusions.
- Other localized musculoskeletal pain conditions.

Dosage and Administration

- How to Use:
 - Apply a thin layer of the gel to the affected area.
 - Gently massage until absorbed.
 - Use 3–4 times daily or as directed by a healthcare professional.
- Precautions for Use:
 - Do not apply to broken, wounded, or irritated skin.

 Avoid contact with eyes, mouth, and mucous membranes.

- Protect from direct sunlight and heat.
- · Keep out of reach of children.

Contraindications

- Hypersensitivity to diclofenac, salicylates, menthol, or any component of the gel.
- Patients with aspirin-sensitive asthma or severe allergic reactions to NSAIDs.

Warnings and Precautions

- For external use only.
- Wash hands thoroughly after application.
- Avoid prolonged use without medical supervision.
- Consult a healthcare provider before use during pregnancy, breastfeeding, or in children under 12 years.

Adverse Effects

- Common: Mild skin irritation, redness, or rash at the application site.
- Rare: Allergic reactions such as itching, swelling, or difficulty breathing (seek medical attention if these occur).

Storage

Store at a temperature below 25°C.

Benefits

- Fast-acting relief from pain and inflammation.
- Non-greasy formula that absorbs quickly.
- Suitable for localized application, reducing systemic side effects compared to oral medications.

Note: This product is intended for shortterm use and symptomatic relief. If pain or inflammation persists or worsens, consult a healthcare professional.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for