

Decplis[®] MR Tablet

Chlorzoxazone (250mg) + Diclofenac (50mg) + Paracetamol (325mg) Tablet

This combination tablet is used for the effective management of pain, inflammation, and muscle spasms. It provides multi-faceted relief by combining a muscle relaxant, an anti-inflammatory agent, and an analgesic.

Composition:

- **Chlorzoxazone: 250 mg**
 - **Diclofenac Sodium: 50 mg**
 - **Paracetamol: 325 mg**
-

Indications:

This medication is indicated for:

1. **Acute Musculoskeletal Pain**
 2. **Inflammatory Conditions with Muscle Spasms**
 3. **Post-Traumatic Pain**
 4. **Arthritis with Associated Muscle Stiffness**
 5. **Back Pain and Neck Pain**
-

Pharmacological Actions:

1. **Chlorzoxazone:** A muscle relaxant that relieves muscle spasms and stiffness.
 2. **Diclofenac:** A non-steroidal anti-inflammatory drug (NSAID) that reduces inflammation and alleviates pain.
 3. **Paracetamol:** An analgesic and antipyretic that provides pain relief and reduces fever.
-

Dosage & Administration:

- **Adults:** One tablet 1–3 times daily, after meals, or as prescribed by the physician.
- **Pediatric Use:** Not recommended unless directed by a healthcare provider.

Administration Instructions:

- Take the tablet with a glass of water after food to minimize gastrointestinal discomfort.
-

Benefits:

- Provides rapid and effective relief from pain and inflammation.
 - Relaxes muscles and improves mobility.
 - Reduces fever and associated discomfort.
-

Side Effects:

- **Common Side Effects:**
 - Nausea or vomiting
 - Drowsiness
 - Stomach upset or acidity
 - **Rare but Serious Side Effects:**
 - Allergic reactions (rash, swelling, or difficulty breathing)
 - Liver toxicity (jaundice, dark urine)
 - Gastrointestinal bleeding
-

Precautions:

- 1. Liver or Kidney Impairment:** Use with caution in individuals with liver or kidney disorders.
 - 2. Alcohol Use:** Avoid alcohol consumption to reduce the risk of liver damage.
 - 3. Pregnancy and Breastfeeding:** Not recommended unless prescribed by a physician.
 - 4. Driving or Operating Machinery:** May cause drowsiness; exercise caution.
-

Drug Interactions:

- **Avoid concurrent use with:**
 - **Other NSAIDs (e.g., ibuprofen, aspirin)** to prevent additive side effects.
 - **Blood thinners like warfarin (increased bleeding risk).**
 - **Alcohol (increased risk of liver damage).**
-

Overdose Symptoms:

- **Symptoms may include severe nausea, vomiting, dizziness, or abdominal pain. Seek immediate medical attention in case of overdose.**
-

Storage:

- **Store in a cool, dry place below 25°C.**
 - **Keep out of reach of children.**
-

Key Counseling Points:

- 1. Follow the prescribed dosage; do not exceed the recommended dose.**
- 2. Take the tablet after meals to reduce the risk of gastric irritation.**
- 3. Avoid activities requiring alertness if drowsiness occurs.**

- 4. Report any signs of liver damage, such as yellowing of the skin or dark urine, to your doctor immediately.**
 - 5. Consult a physician if pain or inflammation persists beyond a few days.**
-

This Chlorzoxazone + Diclofenac + Paracetamol Tablet is an effective solution for managing pain, inflammation, and muscle spasms, ensuring improved mobility and comfort. Use as directed by your healthcare provider.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for