

# Dectyle 50 Injection

## Dectyle 50 Injection: Comprehensive Information

**Generic Name:** Dexamethasone

**Brand Name:** Dectyle 50

**Strength:** 50mg per mL

**Formulation:** Injectable Solution

**Drug Class:** Corticosteroid

## Description

Dectyle 50 Injection contains **Dexamethasone**, a potent synthetic corticosteroid with anti-inflammatory, immunosuppressant, and antiallergic properties. It is commonly used to treat a variety of conditions, including inflammatory, allergic, and autoimmune disorders.

## Indications and Uses

Dectyle 50 Injection is primarily used for the following conditions:

- Inflammatory Conditions:**
  - Rheumatoid arthritis, osteoarthritis, bursitis**, and other inflammatory joint disorders.
  - Inflammatory bowel disease**, such as Crohn's disease and ulcerative colitis.
- Allergic Reactions:**
  - Severe allergic reactions, such as **allergic rhinitis, asthma, and contact dermatitis**.
  - Anaphylaxis (severe allergic reaction) when steroids are required for symptom control.
- Autoimmune Disorders:**
  - Conditions like **lupus, autoimmune hepatitis, and multiple sclerosis** where the immune system attacks the body's tissues.
- Cancers:**

- As part of chemotherapy regimens, particularly in **lymphoma, leukemia, and myeloma**.

- Acute Conditions:**

- Used in emergency settings for shock, adrenal insufficiency, or acute flare-ups of chronic conditions.

- \*\*Preventing and Treating Nausea and Vomiting:**

- Associated with chemotherapy.

## Dosage and Administration

- Route of Administration:**

- Dectyle 50 Injection is administered **intramuscularly (IM) or intravenously (IV)**.

- General Dosage Guidelines:**

- For Inflammatory Conditions:** Initial dosage varies but typically ranges from **4-20mg** per day depending on the severity of the condition. The dosage is then gradually tapered down.
- For Allergic Reactions and Autoimmune Disorders:** The dose can range from **10-50mg** initially, followed by a tapering regimen.
- For Cancer Chemotherapy:** Typically used as part of combination therapy, with a specific dose prescribed by the oncologist.
- Acute Conditions or Emergencies:** A higher initial dose may be used, followed by reduction based on clinical response.

- Administration Notes:**

- The injection site should be rotated to avoid tissue damage or irritation.
- Monitor the patient closely for side effects, especially during long-term treatment.

## Precautions Before Use

## Do not use Dectyle 50 Injection if you:

- Have a known allergy to dexamethasone or any of its components.
- Have a history of **fungal infections, viral infections, or tuberculosis**.
- Have an untreated infection (except for specific medical circumstances).
- Are pregnant or breastfeeding (unless specifically prescribed by a healthcare professional).

## Use cautiously if you have:

- **Cardiovascular disease** (e.g., high blood pressure, heart failure).
- **Peptic ulcer disease** or **gastric ulcers**.
- **Osteoporosis** or **bone fractures**.
- **Diabetes mellitus** (may cause blood sugar levels to rise).
- **Psychiatric disorders**, such as mood swings or depression, may be exacerbated by corticosteroid use.

## Potential Side Effects

### 1. Common Side Effects:

- Weight gain, fluid retention.
- Increased blood pressure or elevated blood glucose levels.
- Mood changes, including irritability and anxiety.
- Gastrointestinal symptoms, such as indigestion or ulcers.

### 2. Serious Side Effects:

- **Osteoporosis** (bone thinning) with long-term use.
- **Cushing's syndrome** (moon face, abdominal obesity, thinning skin).
- **Adrenal insufficiency** (the body's ability to respond to stress is impaired).
- **Infections**, due to immune suppression.
- **Eye problems** (cataracts, glaucoma).

### 3. Rare but Severe Side Effects:

- Severe allergic reactions (rash, swelling, difficulty breathing).
- **Psychosis** or severe mood disorders.
- **Gastrointestinal bleeding** or perforation.

---

## Drug Interactions

### 1. Medications to Avoid or Monitor:

- **Nonsteroidal anti-inflammatory drugs (NSAIDs)**: May increase the risk of gastrointestinal side effects like ulcers.
- **Diuretics (water pills)**: Can increase the risk of electrolyte imbalances.
- **Antidiabetic medications**: Corticosteroids can elevate blood sugar levels, requiring dose adjustments.
- **Vaccines**: Live vaccines may not be effective during treatment with corticosteroids.
- **Anticoagulants**: Dexamethasone may affect blood clotting, requiring careful monitoring when combined with anticoagulants.

---

## Overdose and Emergency Management

### Symptoms of overdose:

- **Excessive weight gain** and fluid retention.
- **Severe high blood pressure** or elevated blood glucose levels.
- **Psychiatric symptoms**, including confusion or agitation.

### Management:

- Immediate medical attention is necessary. Treatment is symptomatic, and the healthcare provider may adjust the dosage or discontinue the medication.

---

## Storage Instructions

- Store at **room temperature** (20-25°C / 68-77°F).
  - Protect from light and moisture.
  - Keep out of the reach of children.
  - **Do not freeze**.
-

## Warnings and Legal Considerations

- **Pregnancy Category C:** Dectyle 50 should only be used during pregnancy if the benefits outweigh the risks. Consult your healthcare provider before use.
  - **Controlled Substance:** Dectyle 50 Injection is not a controlled substance but should be used cautiously and only as prescribed due to its potent effects.
- 

**Disclaimer:** Always consult with a qualified healthcare provider before using Dectyle 50 Injection. This medication should only be used under professional guidance, and any adjustments or changes to treatment should be made in consultation with your doctor

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for