

Diziderm IT Cream

Ofloxacin 0.75% + Ornidazole 2% + Itraconazole 1% + Clobetasole 0.05% + Preservatives Cream

Composition (Per gram of cream):

- **Ofloxacin** – 0.75% w/w (Broad-spectrum antibiotic)
- **Ornidazole** – 2% w/w (Antiprotozoal & Antibacterial)
- **Itraconazole** – 1% w/w (Antifungal)
- **Clobetasole Propionate** – 0.05% w/w (Corticosteroid)
- **Preservatives** – q.s.

Pharmacological Class:

- **Broad-spectrum Antibacterial, Antiprotozoal, Antifungal, and Corticosteroid Combination**

Indications:

This cream is used for the **treatment of mixed bacterial, fungal, and inflammatory skin infections**, including:

- **Fungal & Bacterial Skin Infections**
- **Dermatitis (Eczema, Atopic & Contact Dermatitis)**
- **Tinea Infections (Ringworm, Athlete's Foot, Jock Itch)**
- **Psoriasis (Inflammatory Skin Conditions)**
- **Seborrheic Dermatitis**
- **Secondary Infections of Pre-existing Skin Disorders**

Mechanism of Action:

- **Ofloxacin (Antibiotic):** Inhibits bacterial DNA replication, preventing bacterial growth.
- **Ornidazole (Antiprotozoal & Antibacterial):** Disrupts DNA structure of anaerobic bacteria and protozoa, leading to their destruction.

- **Itraconazole (Antifungal):** Inhibits ergosterol synthesis, essential for fungal cell membrane formation.
- **Clobetasole (Corticosteroid):** Reduces inflammation, redness, and itching by suppressing immune responses.

Dosage & Administration:

- **Apply a thin layer** to the affected area **twice daily** or as prescribed by a physician.
- **Gently massage until absorbed.**
- **Do not apply on open wounds or broken skin** unless advised by a doctor.
- **Avoid prolonged use (>2 weeks) unless directed** by a healthcare provider.

Contraindications:

- Hypersensitivity to any of the ingredients.
- **Not for viral infections** (e.g., herpes, chickenpox).
- **Not for ophthalmic (eye) use.**
- **Avoid in pregnancy & breastfeeding unless prescribed.**

Warnings & Precautions:

- **Long-term use may cause skin thinning (due to Clobetasole).**
- **May cause antibiotic resistance** if misused.
- **Avoid occlusive dressings**, as they increase absorption and side effects.
- **Do not use on the face unless prescribed.**

Side Effects:

Common:

- Burning, redness, itching, or dryness at the application site.
- Skin thinning (with prolonged use).

Rare:

- **Allergic reactions (rash, swelling, difficulty breathing – seek medical help immediately).**

- **Skin discoloration or stretch marks (prolonged steroid use).**

Drug Interactions:

- Avoid using with other **topical steroids, antibiotics, or antifungals** unless advised.
- **Itraconazole** may interact with systemic antifungal treatments.

Storage Instructions:

- Store at **room temperature (below 30°C)**.
- Protect from direct sunlight and moisture.
- Keep out of reach of children.

Conclusion:

This combination cream is **effective in treating mixed infections** involving bacteria, fungi, and inflammation. **Use under medical supervision** to prevent resistance, side effects, and overuse complications.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for