

## Diziderm NM Cream

### Clobetasol 0.05% w/w + Neomycin 0.5% w/w + Miconazole 2% w/w Cream

#### Composition (Per gram of cream):

- **Clobetasol Propionate I.P.** – 0.05% w/w (Corticosteroid)
- **Neomycin Sulphate I.P.** – 0.5% w/w (Antibiotic)
- **Miconazole Nitrate I.P.** – 2% w/w (Antifungal)
- **Cream Base** – q.s.

#### Pharmacological Class:

- **Corticosteroid, Antibiotic, and Antifungal Combination**

#### Indications:

This **triple-action dermatological cream** is used for the treatment of **mixed inflammatory, fungal, and bacterial skin infections**, including:

- **Tinea Infections** (Ringworm, Athlete's Foot, Jock Itch)
- **Eczema & Dermatitis (Atopic, Seborrheic, and Contact Dermatitis)**
- **Bacterial & Fungal Superinfections**
- **Psoriasis (Inflammatory Skin Condition)**
- **Intertrigo (Skin Fold Infections)**

#### Mechanism of Action:

- **Clobetasol Propionate (Corticosteroid):** Reduces redness, swelling, and itching by suppressing inflammation.
- **Neomycin (Antibiotic):** Inhibits bacterial protein synthesis, preventing and treating bacterial skin infections.
- **Miconazole (Antifungal):** Disrupts fungal cell membranes by inhibiting ergosterol synthesis, stopping fungal growth.

#### Dosage & Administration:

- **Apply a thin layer** to the affected area **twice daily** or as directed by a physician.
- **Gently massage until absorbed.**

- **Avoid prolonged use (>2 weeks) unless prescribed** by a doctor.
- **Do not cover with occlusive dressings unless advised.**

#### Contraindications:

- Hypersensitivity to any of the ingredients.
- **Not for viral infections** (e.g., herpes, chickenpox).
- **Not for ophthalmic (eye) use.**
- **Avoid application on deep wounds or large open sores.**
- **Not recommended during pregnancy & breastfeeding unless prescribed.**

#### Warnings & Precautions:

- **Prolonged use may cause skin thinning & steroid-induced side effects.**
- **May lead to antibiotic resistance if overused.**
- **Avoid application near the eyes, nose, or mouth.**
- **Do not use for more than 2 weeks without medical advice.**

#### Side Effects:

##### Common:

- Mild burning, redness, itching, or irritation at the application site.
- Dry or peeling skin.

##### Rare:

- **Allergic reactions (rash, swelling, difficulty breathing – seek medical help immediately).**
- **Skin thinning or discoloration (prolonged steroid use).**
- **Secondary infections due to overuse.**

#### Drug Interactions:

- Avoid using with other **topical corticosteroids, antibiotics, or antifungals** unless prescribed.
- **Neomycin may interact with other aminoglycoside antibiotics** if used systemically.

*Storage Instructions:*

- Store at **room temperature (below 30°C)**.
- Keep away from direct sunlight and moisture.
- **Keep out of reach of children.**

*Conclusion:*

This **broad-spectrum dermatological cream** effectively treats **fungal, bacterial, and inflammatory skin conditions**. However, **prolonged or unnecessary use should be avoided to prevent resistance and side effects. Use under medical supervision for best results.**

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for