

Dobetye Cream

**Calcium Dobesilate Monohydrate (0.5% w/w)
+ Lignocaine Hydrochloride (3% w/w) +
Hydrocortisone Acetate (0.25% w/w) + Zinc
Oxide (5% w/w) Cream**

This combination cream is formulated to provide symptomatic relief and promote healing in conditions like hemorrhoids, anal fissures, and other anorectal disorders. It combines a capillary stabilizer, a local anesthetic, an anti-inflammatory agent, and a soothing protectant to address pain, inflammation, and discomfort.

Composition (per gram of cream):

- **Calcium Dobesilate Monohydrate I.P. 0.5% w/w:** Improves microcirculation, stabilizes capillaries, and reduces swelling.
- **Lignocaine Hydrochloride I.P. 3% w/w:** Acts as a local anesthetic to relieve pain and itching.
- **Hydrocortisone Acetate I.P. 0.25% w/w:** Reduces inflammation, redness, and swelling.
- **Zinc Oxide I.P. 5% w/w:** Acts as a skin protectant and promotes healing by forming a protective barrier.

Uses:

1. **Hemorrhoids (Piles):**
 - **Relieves pain, itching, and swelling associated with internal and external hemorrhoids.**
2. **Anal Fissures:**

- **Promotes healing of tears in the anal canal while reducing pain and inflammation.**
3. **Pruritus Ani:**
 - **Alleviates itching and irritation in the anal region.**
 4. **Post-Operative Care:**
 - **Aids in managing symptoms and promoting recovery after anorectal surgeries.**

Mechanism of Action:

- **Calcium Dobesilate:** Improves blood flow and reduces capillary permeability, thus minimizing swelling.
- **Lignocaine Hydrochloride:** Blocks nerve signals to provide fast-acting pain relief.
- **Hydrocortisone Acetate:** A corticosteroid that suppresses inflammation and immune responses at the site of application.
- **Zinc Oxide:** Forms a protective barrier to prevent irritation and supports natural healing processes.

Directions for Use:

1. **Wash and pat dry the affected area.**
2. **Apply a thin layer of the cream to the affected area 2-3 times daily or as directed by a healthcare provider.**
3. **Wash hands thoroughly after application.**

Side Effects:

- **Common:**

- Mild burning, itching, or irritation at the application site.
 - Rare (serious):
 - Allergic reactions such as rash, swelling, or difficulty breathing (discontinue use and seek medical attention).
-

Precautions:

- 1. External Use Only: Avoid contact with eyes, mouth, or open wounds.**
 - 2. Pregnancy and Lactation: Use under medical supervision during pregnancy or breastfeeding.**
 - 3. Prolonged Use: Avoid extended use without medical guidance to prevent potential side effects of corticosteroids.**
 - 4. Allergy: Discontinue use if signs of an allergic reaction occur.**
-

Storage:

- Store in a cool, dry place below 25°C.
 - Protect from direct sunlight and keep the tube tightly closed.
 - Keep out of reach of children.
-

Note: This medication should be used as prescribed by a healthcare provider. If symptoms persist or worsen, consult a doctor promptly.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for