

Information For the Use of
Etodolac (300mg) Tablet

Edofoam ER 300 Tablet

Brand Name: Edofoam ER 300 Tablet
Dosage Form: Extended Release Tablet
Composition:
Each extended-release tablet contains:
• Etodolac IP 300 mg

Description:

Edofoam ER 300 Tablet contains Etodolac, a non-steroidal anti-inflammatory drug (NSAID) with analgesic and anti-inflammatory properties. It is indicated for the management of pain and inflammatory conditions associated with musculoskeletal disorders. It helps in reducing pain, inflammation, and joint stiffness, thereby improving mobility and daily functioning.

Indications:

Indicated for the treatment of:

- Osteoarthritis
- Rheumatoid arthritis
- Acute musculoskeletal pain
- Post-operative and inflammatory pain conditions

It helps in relieving pain, swelling, and inflammation associated with joint and soft tissue disorders.

Dosage & Administration:

Dosage should be as directed by the physician.
The tablet should be swallowed whole with water, preferably after meals.
Do not crush or chew the extended-release tablet.
Use the lowest effective dose for the shortest duration necessary.

Pharmacological Action:

- Etodolac inhibits cyclooxygenase (COX) enzymes, thereby reducing the synthesis of prostaglandins responsible for pain and inflammation.
- The extended-release formulation provides sustained therapeutic plasma concentrations for prolonged symptom control.

These actions help in effective management of pain and inflammatory conditions.

Side Effects:

Common side effects include nausea, abdominal pain, dyspepsia, headache, dizziness, and diarrhea. Long-term use may increase the risk of gastrointestinal ulceration, bleeding, or renal impairment. Rare hypersensitivity reactions may occur.

Precautions & Warnings:

- Use with caution in patients with gastrointestinal disorders, cardiovascular disease, or renal impairment
 - Avoid prolonged use without medical supervision
 - Monitor renal and hepatic function during long-term therapy
 - Not recommended in patients with active peptic ulcer disease
 - Use during pregnancy only if clearly advised by the physician
-

Drug Interactions:

May interact with anticoagulants, corticosteroids, antihypertensives, and other NSAIDs. Concomitant use with alcohol may increase gastrointestinal irritation risk.

Storage:

**Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.**

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

TM: Trademark Applied for