

Etoricoxib (10mg) + Linseed Oil (30mg) + Menthol (50mg) + Methyl Salicylate (100mg)

Etoscrib Gel Ointment

**Etoricoxib (10 mg) + Linseed Oil (30 mg) +
Menthol (50 mg) + Methyl Salicylate (100 mg)
Ointment**

Composition:

Each gram of ointment contains:

- **Etoricoxib: 10 mg**
- **Linseed Oil: 30 mg**
- **Menthol: 50 mg**
- **Methyl Salicylate: 100 mg**

Pharmacological Class:

- **Etoricoxib: Selective COX-2 Inhibitor (NSAID)**
- **Linseed Oil: Lubricant and anti-inflammatory agent**
- **Menthol: Counterirritant**
- **Methyl Salicylate: Topical analgesic (Salicylate compound)**

Indications:

This ointment is used for the topical treatment of:

- **Pain and inflammation associated with musculoskeletal conditions such as arthritis, sprains, strains, and muscle or joint pain.**
- **Post-traumatic pain (e.g., bruises, sprains, or soft tissue injuries).**
- **Neck, back, and joint pain relief.**
- **Localized pain relief in conditions like tendonitis or bursitis.**

Mechanism of Action:

- **Etoricoxib: Selectively inhibits the COX-2 enzyme, reducing local inflammation and pain.**
- **Linseed Oil: Contains omega-3 fatty acids which have anti-inflammatory properties, promoting skin hydration and nourishment.**

- **Menthol: Acts as a counterirritant by producing a cooling sensation on the skin, which helps alleviate pain by distracting the brain from deeper pain signals.**
- **Methyl Salicylate: A local analgesic that works by reducing inflammation and improving blood circulation to the affected area, helping to reduce pain and discomfort.**

Dosage and Administration:

- **Apply a thin layer of the ointment to the affected area 2-3 times daily, or as directed by a healthcare provider.**
- **Gently massage the ointment into the skin until fully absorbed.**
- **Avoid applying to broken or irritated skin.**
- **Wash hands after applying the ointment, unless the hands are the area being treated.**

Contraindications:

- **Hypersensitivity or allergy to any of the ingredients (Etoricoxib, Linseed Oil, Menthol, or Methyl Salicylate).**
- **History of hypersensitivity to NSAIDs or salicylates.**
- **Not recommended for use in patients with open wounds, dermatitis, or skin infections in the application area.**
- **Avoid use in children under 12 years of age.**
- **Use with caution in patients with a history of gastrointestinal ulcers, asthma, or kidney and liver disease.**

Warnings and Precautions:

- **Gastrointestinal Irritation: Although applied topically, Etoricoxib may still have systemic absorption, and long-term or excessive use could lead to gastrointestinal discomfort or bleeding.**

- **Skin Reactions:** Discontinue use if skin irritation, rash, or allergic reactions (such as swelling, redness, or itching) occur.
- **Pregnancy and Breastfeeding:**
 - **Pregnancy Category C:** Should be avoided during the third trimester due to potential risks to the fetus.
 - **Breastfeeding:** Consult a healthcare provider before use, as Etoricoxib may pass into breast milk in small amounts.
- **Application Near Eyes or Mucous Membranes:** Avoid contact with the eyes, mucous membranes, and sensitive areas. If contact occurs, rinse thoroughly with water.

Side Effects:

Common side effects:

- Skin irritation (e.g., redness, itching, rash)
- Burning or tingling sensation at the application site
- Allergic reactions (e.g., swelling, hives)

Rare but serious side effects:

- Severe allergic reactions (anaphylaxis, difficulty breathing)
- Dermatitis or severe skin reactions
- Persistent or worsening pain or inflammation after use

Drug Interactions:

- **Topical corticosteroids or other topical analgesics:** Concurrent use may increase the risk of skin irritation or other adverse reactions.
- **Oral NSAIDs (e.g., ibuprofen, aspirin):** Caution should be used when combining oral NSAIDs with this topical treatment due to the potential for additive systemic effects.
- **Anticoagulants (e.g., warfarin):** Monitor for any signs of bleeding as Methyl Salicylate has blood-thinning properties.

Storage:

- Store at room temperature (below 25°C) in a cool, dry place, away from heat and direct sunlight.
- Keep the container tightly closed and out of reach of children.

Packaging:

- Available in tubes of 30 g, 50 g, or 100 g.

Manufacturer:

- [Manufacturer Name]

Note:

This ointment is intended for external use only. For best results, use as directed by your healthcare provider and avoid excessive or prolonged use. If symptoms persist or worsen, consult your healthcare provider.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for