

Information For the Use of
Fluorometholone IP 0.1% w/v + Benzalkonium Chloride solution IP 0.02% v/v Eye Drops

Flurmine Eye Drops

Brand Name: Flurmine Eye Drop

Dosage Form: Eye Drops

Composition:

Each ml contains:

- Fluorometholone IP 0.1% w/v
- Benzalkonium Chloride Solution IP 0.02% v/v (as preservative)
- Sterile ophthalmic vehicle q.s.

Description:

Flurmine Eye Drop contains Fluorometholone, a corticosteroid indicated for the treatment of steroid-responsive inflammatory conditions of the eye. It helps in reducing ocular inflammation, redness, swelling, and discomfort associated with allergic and inflammatory eye disorders. The formulation provides localized anti-inflammatory action with minimal systemic absorption when used as directed.

Indications:

Indicated for the treatment of:

- Steroid-responsive ocular inflammation
- Allergic conjunctivitis
- Anterior segment inflammatory conditions of the eye
- Post-operative ocular inflammation as advised by the physician

It helps in relieving redness, swelling, irritation, and inflammatory symptoms affecting the eye.

Dosage & Administration:

Instill the prescribed number of drops into the affected eye(s) as directed by the physician.

Wash hands thoroughly before administration.

Avoid touching the dropper tip to any surface to prevent contamination.

If using multiple ophthalmic medications, maintain an interval of at least 5–10 minutes between applications.

Pharmacological Action:

- Fluorometholone is a corticosteroid that suppresses inflammatory responses by inhibiting the release of inflammatory mediators and reducing leukocyte infiltration.
- It decreases capillary permeability, edema, redness, and tissue inflammation within ocular tissues.

These actions help in effective management of inflammatory eye conditions.

Side Effects:

Common side effects include transient burning, stinging, blurred vision, eye irritation, and discomfort after instillation.

Prolonged use may increase intraocular pressure, delay wound healing, or increase the risk of secondary ocular infections.

Rarely, cataract formation may occur with long-term use.

Precautions & Warnings:

- For ophthalmic use only
- Prolonged use may lead to increased intraocular pressure or glaucoma
- Monitor intraocular pressure during extended therapy
- Use with caution in patients with ocular infections or a history of glaucoma
- Remove contact lenses before administration and reinsert only as advised by the physician
- Do not use for longer than prescribed without ophthalmic evaluation

Drug Interactions:

No significant systemic drug interactions are expected due to minimal systemic absorption.
Concurrent use with other ophthalmic corticosteroids may increase the risk of adverse ocular effects.

Storage:

Store below 25°C in a cool, dry place.

Protect from light.

Do not freeze.

Keep the bottle tightly closed after use.

Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113

TM: Trademark Applied for