

Flupitale P Tablet

Flupirtine (100 mg) + Paracetamol (325 mg) Tablet Information

Composition:

Each tablet contains:

- Flupirtine maleate: 100 mg
 - Paracetamol (Acetaminophen): 325 mg
-

Description

This combination tablet is used for managing pain and inflammation. Flupirtine is a centrally acting non-opioid analgesic with muscle-relaxing properties, while paracetamol is an antipyretic and analgesic that helps reduce pain and fever. Together, they provide enhanced pain relief, especially for musculoskeletal and neuropathic pain.

Indications

This medication is prescribed for:

- Acute and chronic pain management (e.g., back pain, musculoskeletal pain).
 - Neuropathic pain.
 - Post-operative pain.
 - Pain associated with conditions like osteoarthritis or fibromyalgia.
-

Mechanism of Action

1. **Flupirtine:** Acts as a selective neuronal potassium channel opener (SNEPCO), which helps in pain relief by stabilizing nerve cells and reducing overexcitation. It also exhibits muscle-relaxing effects.
2. **Paracetamol:** Inhibits prostaglandin synthesis in the central nervous system, alleviating pain and lowering fever.

Dosage and Administration

- **Typical Dosage:** One tablet 2-3 times daily, or as prescribed by a healthcare professional.
 - **Route:** Oral, with or without food.
 - **Duration:** Use only as directed by a healthcare provider to avoid long-term side effects.
-

Contraindications

- Hypersensitivity to flupirtine, paracetamol, or any excipients.
 - Severe liver impairment or active liver disease.
 - Alcohol dependence.
 - Patients with a history of encephalopathy.
-

Precautions and Warnings

- **Hepatotoxicity:** Monitor liver function during prolonged use, as both components can affect the liver.
 - **Kidney Function:** Caution in patients with renal impairment.
 - Avoid excessive alcohol consumption during treatment.
 - Not recommended for pediatric use.
-

Side Effects

Common side effects include:

- Drowsiness
- Dizziness
- Nausea or vomiting
- Fatigue
- Abdominal discomfort

Rare but severe side effects include:

- Hepatotoxicity (elevated liver enzymes, jaundice)
 - Allergic reactions (rash, itching, swelling)
-

Drug Interactions

- **Alcohol:** Increased risk of liver damage.
 - **Other hepatotoxic drugs:** Monitor liver function closely.
 - **CNS depressants:** Enhanced sedative effects with flupirtine.
-

Storage

- Store at room temperature (20-25°C), away from direct sunlight and moisture.
 - Keep out of reach of children.
-

Note: This medication should only be used under medical supervision. Always consult your doctor before starting or stopping the treatment.

Disclaimer:

This medication should be used under medical supervision. Consult your healthcare provider for appropriate use, especially if symptoms persist or worsen.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for