

Levonoble Syrup

Levocetirizine (2.5 mg/5 ml) Suspension

Description:

Levocetirizine is a second-generation antihistamine that blocks the effects of histamine, a substance produced by the body during allergic reactions. Levocetirizine helps relieve common allergy symptoms such as sneezing, runny nose, itchy or watery eyes, and itching. This oral suspension form is specifically designed for children or individuals who have difficulty swallowing tablets. It is also used to treat chronic urticaria (hives) and symptoms related to allergic rhinitis.

Composition:

Each 5 ml of suspension contains:

- Levocetirizine: 2.5 mg

Indications:

Levocetirizine 2.5 mg/5 ml suspension is indicated for:

- **Allergic Rhinitis:**
 - Seasonal allergic rhinitis (hay fever).
 - Perennial allergic rhinitis (e.g., allergies to dust, mold, pet dander).
- **Chronic Urticaria (Hives):** For relief of symptoms like itching, rash, and skin discomfort caused by chronic urticaria.

Dosage and Administration:

Adults and Children Over 6 Years:

- The usual dose is 10 ml (5 mg) once daily, preferably taken in the evening.
Children (2 to 6 years):
- The recommended dose is 5 ml (2.5 mg) once daily, or as prescribed by a healthcare provider.
- **Children Under 2 Years:** Consult a healthcare provider for appropriate dosing for infants under 2 years of age.

Contraindications:

Levocetirizine 2.5 mg/5 ml suspension is contraindicated in:

- Hypersensitivity to Levocetirizine, cetirizine, or any of the ingredients of the suspension.
- **Severe Renal Impairment:** Patients with severely reduced kidney function (creatinine clearance less than 10 mL/min) should avoid using this medication unless directed by a healthcare provider.
- **Pregnancy and Breastfeeding:** Levocetirizine should be used during pregnancy only if the benefits outweigh the risks, and it should be avoided during breastfeeding unless prescribed by a healthcare provider.

Warnings and Precautions:

- **Drowsiness:** Although Levocetirizine is less likely to cause drowsiness compared to first-generation antihistamines, some individuals may still experience mild sedation. It's recommended to avoid activities requiring full alertness, such as driving or operating machinery, until personal tolerance to the medication is known.

- **Renal Impairment:** Dose adjustments may be necessary for patients with renal impairment (e.g., reduced kidney function). Consult a healthcare provider for proper dosing if kidney function is compromised.
- **Pregnancy and Breastfeeding:** Levocetirizine should be used during pregnancy only if the potential benefits justify the risks to the fetus. It is excreted in breast milk, so consult a healthcare provider before using while breastfeeding.
- **Children under 2 years:** The safety and efficacy of Levocetirizine suspension in children under 2 years have not been established.

- Store at room temperature (between 15°C to 30°C), away from moisture and heat.
- Keep the bottle tightly closed and out of reach of children.
- Do not use the suspension after the expiration date printed on the packaging.
- Shake well before use to ensure even distribution of the active ingredient.

Note:

Levocetirizine 2.5 mg/5 ml suspension is a prescription-only medication. Use it only as directed by a healthcare provider.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for

Side Effects:

- **Common Side Effects:**
 - Drowsiness (rare)
 - Headache
 - Dry mouth
 - Nausea or mild gastrointestinal discomfort
 - Fatigue or dizziness
 - Sore throat
- **Less Common Side Effects:**
 - Abdominal pain or discomfort
 - Skin rash or itching
- **Serious Side Effects:**
 - **Severe allergic reactions:** Symptoms like difficulty breathing, swelling of the face, lips, tongue, or throat, and hives.
 - **Kidney or liver issues:** Symptoms such as yellowing of the skin or eyes (jaundice), dark urine, or unusual fatigue.
 - **Severe drowsiness or dizziness:** If you experience excessive sedation, consult a healthcare provider for further evaluation.

Storage: