

Information For the Use of
Lornoxicam (8mg) + Paracetamol (325mg) Tablet

Loxifrit P Tablet

Brand Name: Loxifrit P Tablet

Dosage Form: Tablet

Composition:

Each film-coated tablet contains:

- Lornoxicam IP 8 mg
- Paracetamol IP 325 mg

Description:

Loxifrit P Tablet is a fixed-dose combination of Lornoxicam, a non-steroidal anti-inflammatory drug (NSAID), and Paracetamol, an analgesic and antipyretic agent. It is indicated for the symptomatic relief of pain and inflammation associated with musculoskeletal and inflammatory disorders. It helps in reducing pain, inflammation, and fever, thereby improving mobility and patient comfort.

Indications:

Indicated for the treatment of:

- Acute musculoskeletal pain
- Osteoarthritis and rheumatoid arthritis
- Post-operative pain
- Dental pain
- Low back pain
- Sprains, strains, and soft tissue injuries
- Fever associated with painful inflammatory conditions

It helps in relieving pain, reducing inflammation, and improving functional mobility.

Dosage & Administration:

Dosage should be as directed by the physician.

The tablet should preferably be taken after meals with sufficient water to minimize gastrointestinal irritation.

Do not exceed the recommended dosage or duration of treatment.

Pharmacological Action:

- Lornoxicam inhibits cyclooxygenase (COX-1 and COX-2) enzymes, thereby reducing the synthesis of prostaglandins responsible for pain and inflammation.
- Paracetamol exerts analgesic and antipyretic effects by inhibiting prostaglandin synthesis primarily within the central nervous system and regulating the hypothalamic heat-regulating center.

Together, these mechanisms provide effective relief from pain, inflammation, and fever.

Side Effects:

Common side effects include nausea, abdominal discomfort, dyspepsia, headache, dizziness, vomiting, and diarrhea.

Prolonged or excessive use may increase the risk of gastrointestinal bleeding, hepatic impairment, renal dysfunction, or hypersensitivity reactions.

Precautions & Warnings:

- Use with caution in patients with a history of peptic ulcer disease or gastrointestinal bleeding
- Monitor liver and renal function during prolonged therapy
- Avoid concomitant use with other NSAIDs or Paracetamol-containing products
- Use cautiously in elderly patients and those with hepatic, renal, or cardiovascular impairment
- Not recommended during the third trimester of pregnancy unless clearly indicated by the physician

Drug Interactions:

May interact with anticoagulants, antiplatelet agents, corticosteroids, diuretics, antihypertensive agents, methotrexate, and other NSAIDs.

Concurrent use with alcohol may increase the risk of hepatotoxicity and gastrointestinal adverse effects.

Storage:

Store below 25°C in a cool, dry place.

Protect from light and moisture.

Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,

Panchkula, Haryana -134113

TM: Trademark Applied for