# **Megendo 40 SR Tablet**

# Megestrol Acetate 40 mg SR Tablet

# **Generic Name:**

Megestrol Acetate

# **Strength:**

40 mg

# **Formulation:**

Sustained Release (SR) tablet

# **Uses:**

Megestrol Acetate 40 mg SR is primarily used in the treatment of the following conditions:

### 1. Anorexia and Cachexia:

 Used to stimulate appetite and promote weight gain in patients suffering from conditions such as cancer or AIDS. This medication helps improve nutritional intake and body mass in those with severe unintentional weight loss.

# 2. Breast Cancer (Adjuvant Therapy):

 Used in certain cases to treat breast cancer, particularly in postmenopausal women with hormone receptor-positive tumors, by lowering estrogen levels and helping to slow the growth of hormone-dependent cancer cells.

# 3. Endometrial Cancer:

 May be used for the treatment of endometrial cancer by inhibiting the growth of estrogen-dependent tumor cells.

# **Mechanism of Action:**

Megestrol acetate is a **synthetic progestin** (a type of synthetic progesterone). It exerts its effects through the following actions:

- Appetite Stimulation: Increases appetite and food intake by acting on the hypothalamus, the part of the brain responsible for regulating hunger.
- Anti-estrogenic Effect: It can block the action of estrogen on certain tumors (such as in breast and endometrial cancer), slowing their growth by modifying hormone levels.
- Inhibition of Tumor Growth: Helps reduce the growth of hormone-sensitive cancers by altering hormonal balance in the body.

# **Dosage:**

- Typical Dose for Anorexia and Cachexia:
  - o 40 mg SR tablet, taken once daily.
  - The dosage and duration will depend on the patient's condition and the response to treatment.
- Typical Dose for Cancer Treatment:
  - The dose for cancer treatment may vary based on the specific type of cancer and the physician's recommendations.
- Administration:
  - Swallow the sustained-release tablet whole, without chewing or crushing it.
  - o It can be taken with or without food.
  - It is recommended to take it at the same time each day for consistency.

# **Precautions:**

- Pregnancy and Breastfeeding:
  - Megestrol acetate is contraindicated during pregnancy. It should not be used by pregnant women due to potential risks to the fetus.
  - It is also not recommended for breastfeeding mothers unless advised by a healthcare provider.
  - Liver Disease:

 Caution is required in patients with liver disease, and the medication should be used under close monitoring.

# Blood Clotting Disorders:

 Megestrol acetate may increase the risk of blood clots, especially in patients with a history of blood clots or stroke.

#### Diabetes:

 The drug may affect blood sugar levels, so people with diabetes should monitor their blood sugar carefully while on this medication.

# **Side Effects:**

# • Common:

- Weight gain
- Increased appetite
- o Fluid retention
- Swelling in the legs, feet, or abdomen
- Hot flashes or night sweats

### • Less Common:

- Nausea or vomiting
- Abdominal discomfort
- Fatigue or tiredness
- Depression or mood changes

# • Serious (Seek Medical Attention):

- Severe allergic reactions (rash, itching, swelling of the face or throat, difficulty breathing)
- Symptoms of blood clots (pain, redness, or swelling in the legs, sudden chest pain, or difficulty breathing)
- Severe abdominal pain or jaundice (yellowing of the skin or eyes)

# **Drug Interactions:**

Megestrol acetate may interact with the following medications:

# Anticoagulants (e.g., warfarin):

 May increase the risk of bleeding. Regular monitoring of blood clotting parameters is advised.

# Diabetes Medications:

 Megestrol acetate can interfere with blood sugar control, so patients with diabetes should closely monitor their blood glucose levels.

# • Hormonal Treatments:

 Use with caution in patients taking other hormonal treatments, such as hormone replacement therapy or birth control.

### Corticosteroids:

 Combining megestrol with corticosteroids may increase the risk of **fluid retention** and other related side effects.

Always inform your healthcare provider of all medications, supplements, and herbal products you are taking.

# **Storage:**

- Store at 20°C to 25°C (68°F to 77°F).
- Keep the tablets in a dry place, protected from light and moisture.
- Keep out of reach of children.

# Note:

- Megestrol acetate is **not a cure** for cancer but is used to manage symptoms, improve appetite, and help patients maintain body weight.
- It should only be used under the supervision of a healthcare provider, and the dosage should be followed strictly as prescribed.
- Regular monitoring is important during treatment to check for any adverse effects, especially in patients with pre-existing conditions such as diabetes or cardiovascular disease.
- Do not adjust or stop the medication without consulting your healthcare provider.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for