

Information For the User  
Mirabegron (50mg) + Solifenacin (5mg) Tablet

# Miratyle S 50/5 Tablet

**Composition:**

Each film-coated tablet contains:

**Mirabegron** ..... 50 mg  
**Solifenacin Succinate** ..... equivalent to  
Solifenacin 5 mg  
Excipients ..... q.s.

**Description:**

Miratyle S 50/5 Tablet is a fixed-dose combination used in the management of overactive bladder (OAB) with symptoms such as urgency, increased urinary frequency, and urge incontinence. Mirabegron is a  $\beta_3$ -adrenoceptor agonist that relaxes the detrusor smooth muscle during the storage phase, increasing bladder capacity. Solifenacin is an antimuscarinic agent that reduces involuntary bladder contractions by blocking M3 muscarinic receptors.

**Indications:**

Overactive bladder with urinary urgency and frequency  
Urge urinary incontinence  
Neurogenic detrusor overactivity (as advised by physician)

**Mechanism of Action:**

Mirabegron activates  $\beta_3$ -adrenoceptors in the bladder wall, leading to relaxation of the detrusor muscle and increased bladder capacity. Solifenacin blocks muscarinic M3 receptors in the bladder, decreasing detrusor overactivity and improving continence.

**Dosage and Administration:**

One tablet once daily, swallowed whole with water  
May be taken with or without food  
Dose adjustment may be required in renal or hepatic impairment

**Side Effects:**

Dry mouth, constipation  
Urinary tract infection  
Headache, dizziness, palpitations  
Increased blood pressure

**Precautions:**

Monitor blood pressure regularly during treatment  
Use with caution in patients with severe renal or hepatic impairment  
Avoid in patients with uncontrolled hypertension  
Assess for urinary retention before initiating therapy

**Storage:**

Store in a cool, dry place below 25°C

Protect from light and moisture

Keep out of reach of children

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for