Moxyplit 250 LB Capsule

Amoxycillin I.P. 250mg + Lactic Acid Bacillus (1.66 Million CFU) Capsule

This combination capsule is designed to treat bacterial infections effectively while minimizing gastrointestinal side effects.

Amoxycillin acts as a potent antibiotic, and Lactic Acid Bacillus helps maintain gut flora during antibiotic therapy, reducing the risk of diarrhea and other digestive disturbances.

Amoxycillin: Targets bacterial cell walls, causing cell death and resolving infections.

 Lactic Acid Bacillus: Produces lactic acid, creating an environment unfavorable for harmful bacteria while supporting the growth of beneficial microorganisms in the gut.

Composition:

- Amoxycillin I.P. 250mg: A broadspectrum penicillin-class antibiotic that inhibits bacterial growth by disrupting cell wall synthesis.
- Lactic Acid Bacillus (1.66 Million CFU):
 A probiotic that helps restore the natural balance of intestinal bacteria disrupted by antibiotics.

Dosage & Administration:

- Recommended Dose:
 - Adults and children as directed by a physician, typically one capsule two to three times a day.
- Route of Administration: Oral, taken with or after meals.
- Duration: Complete the full course of therapy as prescribed to ensure effective treatment and avoid antibiotic resistance.

Uses:

- 1. Bacterial Infections:
 - Respiratory tract infections (e.g., sinusitis, bronchitis, pneumonia)
 - o Ear infections (otitis media)
 - Urinary tract infections (UTIs)
 - Skin and soft tissue infections
- 2. Prevention of Antibiotic-Associated Gastrointestinal Side Effects:
 - Maintains healthy gut flora to prevent diarrhea and dysbiosis caused by antibiotics.

Contraindications:

- Hypersensitivity or allergy to Amoxycillin, penicillins, or any capsule ingredients.
- History of severe allergic reactions to beta-lactam antibiotics.

Side Effects:

- Common:
 - o Nausea, vomiting, or diarrhea
 - Abdominal discomfort
 - Mild allergic reactions (e.g., rash, itching)
- Rare:

Mechanism of Action:

- Severe allergic reactions (anaphylaxis)
- Clostridioides difficile-associated diarrhea

Precautions:

- Inform your doctor about any known allergies, especially to antibiotics.
- Use with caution in individuals with kidney or liver impairment.
- Avoid skipping doses to prevent the development of antibiotic resistance.
- Alcohol consumption may exacerbate side effects and should be avoided during treatment.

Storage:

- Store in a cool, dry place below 25°C.
- Keep away from direct sunlight and moisture.
- Keep out of reach of children.

Note: This medication is effective only against bacterial infections and will not work for viral illnesses like the common cold. Always use this medicine under medical supervision. Consult your doctor if side effects persist or worsen.

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for