

Ofloglee OZ Tablet

**Ofloxacin (200 mg) + Ornidazole (500 mg)
Tablet**

Description:

This combination tablet contains Ofloxacin, a fluoroquinolone antibiotic, and Ornidazole, an anti-protozoal and anti-bacterial agent. Together, they provide broad-spectrum coverage against bacterial and protozoal infections, making it highly effective in treating mixed infections.

Composition:

- **Ofloxacin: 200 mg**
 - **Ornidazole: 500 mg**
-

Indications:

This medication is indicated for the treatment of:

1. **Mixed Bacterial and Protozoal Infections**
 - Diarrhea and dysentery of bacterial and protozoal origin.
 - Giardiasis and amoebiasis.
2. **Gynecological Infections: Pelvic inflammatory disease.**
3. **Abdominal Infections: Peritonitis and intra-abdominal abscesses.**
4. **Dental Infections: Periodontitis and other oral cavity infections.**
5. **Post-surgical Infections: Particularly in the gastrointestinal or genitourinary tract.**

Mechanism of Action:

1. **Ofloxacin: Inhibits bacterial DNA gyrase and topoisomerase IV, essential for bacterial DNA replication, transcription, and repair, thereby killing bacteria.**
 2. **Ornidazole: Disrupts the DNA of anaerobic bacteria and protozoa, inhibiting their growth and reproduction.**
-

Dosage and Administration:

- **Usual Adult Dose: One tablet twice daily or as directed by a physician.**
 - **Route: Oral, taken with or after meals to avoid gastrointestinal discomfort.**
-

Side Effects:

- **Common: Nausea, vomiting, diarrhea, dizziness, headache, or metallic taste.**
 - **Rare but Serious: Allergic reactions, tendon rupture, or neurological symptoms like seizures.**
-

Precautions:

1. **Avoid alcohol during treatment as Ornidazole may cause a disulfiram-like reaction (nausea, vomiting, and flushing).**

- 2. Use cautiously in patients with liver or kidney impairment, epilepsy, or neurological disorders.**
 - 3. Not recommended for children, pregnant, or breastfeeding women unless prescribed by a doctor.**
 - 4. Avoid taking antacids, calcium, or iron supplements within 2 hours of this medication as they may interfere with absorption.**
-

Contraindications:

- Hypersensitivity to fluoroquinolones, nitroimidazole derivatives, or any component of the tablet.**
-

Storage:

- Store in a cool, dry place, away from direct sunlight and moisture.**
 - Keep out of reach of children.**
-

Note: This medication should be taken only under medical supervision. It is important to complete the prescribed course to prevent the development of drug resistance.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for