

Information For the User

Ofloxacin (0.3% w/v) + Dexamethasone (0.1% w/v) + Hydroxypropylmethylcellulose (0.25% w/v)
Eye/Ear Drop

Ofloglee DH Eye/Ear Drop

Composition:

Each ml contains:

Ofloxacin I.P. 0.3% w/v

Dexamethasone Sodium Phosphate I.P. eq. to Dexamethasone 0.1% w/v

Hydroxypropylmethylcellulose 0.25% w/v

Benzalkonium Chloride Solution I.P. ... 0.02% w/v (as preservative)

Sterile aqueous base q.s.

Description:

Ofloglee DH Eye/Ear Drop is a sterile combination of a broad-spectrum fluoroquinolone antibiotic (Ofloxacin), an anti-inflammatory corticosteroid (Dexamethasone), and a lubricant (Hydroxypropylmethylcellulose). It is indicated for the treatment and symptomatic relief of bacterial infections of the eye and ear, associated with inflammation, redness, and discomfort. The formulation also provides ocular lubrication to relieve dryness and irritation.

Indications:

Bacterial conjunctivitis, keratitis, blepharitis
Post-operative ocular inflammation and infection prophylaxis
Otitis externa and otitis media with intact tympanic membrane
Relief of ocular irritation and dryness

Mechanism of Action:

Ofloxacin inhibits bacterial DNA gyrase and topoisomerase IV, preventing bacterial DNA replication and transcription.
Dexamethasone suppresses inflammatory mediators, reducing swelling, redness, and pain.
Hydroxypropylmethylcellulose acts as a lubricant, maintaining tear film stability and protecting corneal epithelium.

Dosage and Administration:

Eye: Instill 1–2 drops into the affected eye(s) every 4–6 hours, or as directed by the physician.

Ear: Instill 2–3 drops into the affected ear(s) twice or thrice daily, as prescribed.

Use aseptically. Do not touch the dropper tip to any surface.

Side Effects:

Transient burning or stinging sensation
Blurred vision
Itching, redness, or swelling
Prolonged use may increase intraocular pressure or risk of secondary infections

Precautions:

Avoid long-term use without medical supervision
Do not use in viral or fungal ocular/aural infections
Caution in patients with glaucoma

Contact lens wearers should remove lenses before instillation and wait at least 15 minutes before reinsertion

Storage:

Store below 25°C in a dry, dark place. Protect from direct light. Keep the container tightly closed and out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for