

# Ondotyle 8 MD Tablet

## Ondansetron (8 mg) MD Tablet

---

### Description:

Ondansetron 8 mg MD (mouth-dissolving) tablet is an antiemetic medication used to prevent and treat nausea and vomiting caused by various conditions. It belongs to the 5-HT<sub>3</sub> receptor antagonist class, which works by blocking serotonin in the brain and gastrointestinal tract, reducing the sensation of nausea.

---

### Composition:

- Ondansetron: 8 mg
- 

### Indications:

This medication is indicated for the prevention and treatment of:

1. Chemotherapy-Induced Nausea and Vomiting (CINV).
  2. Radiotherapy-Induced Nausea and Vomiting.
  3. Post-Operative Nausea and Vomiting (PONV).
  4. Nausea and Vomiting Due to Gastrointestinal Conditions.
- 

### Mechanism of Action:

Ondansetron works by selectively blocking 5-HT<sub>3</sub> receptors in the brain and gastrointestinal tract, preventing serotonin from triggering the vomiting reflex.

---

### Dosage and Administration:

- Usual Adult Dose:
    - One 8 mg MD tablet taken 30 minutes before chemotherapy, radiotherapy, or surgery.
    - Repeat every 8–12 hours as needed or as directed by a physician.
  - Route: Place the tablet on the tongue and allow it to dissolve completely; no need for water.
- 

### Side Effects:

- Common: Headache, constipation, diarrhea, or dizziness.
  - Rare but Serious: Allergic reactions, irregular heartbeat, or serotonin syndrome when used with other serotonergic drugs.
- 

### Precautions:

1. Use with caution in patients with:
    - Liver impairment.
    - A history of cardiac conditions (e.g., QT prolongation or arrhythmias).
  2. Avoid use with apomorphine as it may cause severe hypotension and unconsciousness.
  3. Monitor for signs of serotonin syndrome if used with other serotonergic medications.
- 

### Contraindications:

- **Hypersensitivity to Ondansetron or any components of the formulation.**
  - **Congenital long QT syndrome.**
- 

**Storage:**

- **Store in a cool, dry place at room temperature, away from direct sunlight and moisture.**
  - **Keep out of reach of children.**
- 

**Note: Ondansetron MD tablets should only be taken under medical supervision. Follow your doctor's instructions for dosage and duration to ensure effective treatment.**

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for