

Panpital CT Capsule

**Cinitapride (3mg) + Pantoprazole (40mg)
Capsule**

Cinitapride and Pantoprazole are combined in this capsule to provide an effective treatment for gastrointestinal disorders. This combination is used to relieve symptoms associated with conditions like acid reflux, indigestion, and gastric ulcers.

Composition per Capsule:

- Cinitapride: 3 mg
 - Pantoprazole: 40 mg
-

Indications:

- 1. Gastroesophageal Reflux Disease (GERD):**
 - Used to treat symptoms of GERD, including heartburn, acid regurgitation, and discomfort caused by acid reflux.
 - 2. Peptic Ulcers:**
 - Helps in the healing of gastric and duodenal ulcers by reducing stomach acid production and improving gastric motility.
 - 3. Dyspepsia (Indigestion):**
 - Relief from symptoms of indigestion, such as bloating, discomfort, and early satiety.
 - 4. Gastritis:**
 - Treatment for inflammation of the stomach lining, often caused by excess stomach acid.
 - infection, which can lead to peptic ulcers.
-

Dosage & Administration:

- **Adults:**
 - For GERD and Dyspepsia: 1 capsule (3 mg Cinitapride + 40 mg Pantoprazole) once daily, usually before a meal.
 - For Peptic Ulcers or Gastritis: The typical dosage is 1 capsule per day, but treatment duration may vary based on the severity of the condition.

Administration:

- Take the capsule whole, without crushing or chewing.
 - Take it before meals, preferably in the morning.
 - Swallow the capsule with a full glass of water.
-

Benefits of Cinitapride + Pantoprazole Capsule:

- 1. Dual Action:**
 - Cinitapride improves gastric motility, helping with symptoms like bloating and delayed stomach emptying.
 - 2. Prevention of Ulcers:**
 - Reduces stomach acid, thus preventing the formation of new ulcers and aiding in the healing of existing ones.
-

Side Effects:

- **Common Side Effects:**
 - Headache

- Diarrhea or constipation
- Nausea or vomiting
- **Less Common Side Effects:**
 - Flatulence (gas)
 - Dry mouth or altered taste
- **Serious Side Effects (Rare):**
 - Allergic reactions such as difficulty breathing, rash, or swelling
 - Persistent stomach pain or severe nausea

If any of these serious side effects occur, discontinue the medication and seek medical help immediately.

Precautions:

1. **Liver Disease:**
 - Use with caution in patients with liver disease. Dose adjustments may be necessary, particularly for Pantoprazole.
 2. **Kidney Disease:**
 - Caution is required in patients with kidney impairment. Monitoring may be necessary, especially with prolonged use.
 3. **Pregnancy and Breastfeeding:**
 - Consult a doctor before using during pregnancy, as both Cinitapride and Pantoprazole should only be used if absolutely necessary.
-

Drug Interactions:

1. **Antacids or Antireflux Medications:**
 - Combining with other antacid medications may reduce the effectiveness of Pantoprazole.
2. **Antibiotics (e.g., Amoxicillin, Clarithromycin):**
 - These drugs are often used in combination with Pantoprazole for the treatment of *H. pylori*

infection. Caution is advised to avoid drug-drug interactions.

3. Clopidogrel:

- Pantoprazole may reduce the effectiveness of clopidogrel, a blood thinner, so this combination should be avoided or monitored closely.

4. Methotrexate:

- Pantoprazole may increase blood levels of methotrexate, a chemotherapy drug, leading to an increased risk of side effects.
-

Overdose Symptoms:

- Symptoms of overdose may include:
 - Severe dizziness
 - Rapid heartbeat
 - Tremors or muscle twitching

If overdose is suspected, seek immediate medical attention.

Storage:

- Store at room temperature (15–30°C).
- Keep the capsule in a tightly sealed container.

Conclusion:

The Cinitapride (3mg) + Pantoprazole (40mg) Capsule is a highly effective combination for treating a variety of gastrointestinal issues, especially GERD, dyspepsia, and peptic ulcers.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for