

Paraplom Dom Tablet

Domperidone I.P. 10 mg + Paracetamol I.P. 325 mg Tablet

Description:

This combination tablet contains **Domperidone** and **Paracetamol**, formulated to provide relief from pain and nausea or vomiting associated with various conditions, including gastrointestinal disorders and migraines.

Composition:

Each tablet contains:

- **Domperidone I.P.** 10 mg
- **Paracetamol I.P.** 325 mg
- **Excipients:** [List excipients based on formulation].

Pharmacological Actions:

1. **Domperidone:**
 - A dopamine receptor antagonist with prokinetic and antiemetic properties. It enhances gastrointestinal motility and prevents nausea and vomiting by acting on the chemoreceptor trigger zone.
2. **Paracetamol:**
 - An analgesic and antipyretic that works by inhibiting prostaglandin synthesis, thereby reducing pain and fever.

Indications:

This combination tablet is indicated for the symptomatic treatment of:

- Headaches and migraines with associated nausea or vomiting
- Pain and fever accompanied by nausea or vomiting
- Gastric motility disorders with mild to moderate pain

Dosage and Administration:

- **Adults and adolescents (12 years and above):**
One tablet 2-3 times a day, as prescribed by the physician.
 - Maximum daily dose: Do not exceed 3 tablets within 24 hours.
- **Method of Administration:** Swallow the tablet whole with water, preferably before meals to enhance absorption.

Contraindications:

- Hypersensitivity to Domperidone, Paracetamol, or any excipients.
- Severe hepatic impairment.
- Gastrointestinal perforation or obstruction.
- Prolactinoma (tumors affecting the pituitary gland).

Warnings and Precautions:

- Prolonged use of paracetamol may cause liver damage. Use cautiously in patients with hepatic impairment or chronic alcohol consumption.
- Domperidone may cause QT interval prolongation; use with caution in patients with cardiac conditions.
- Avoid exceeding the recommended dose to prevent toxicity.

Side Effects:

Common side effects include:

- Drowsiness
- Dry mouth
- Abdominal cramps

Rare but serious side effects:

- Allergic reactions, including rash or angioedema
- Hepatic dysfunction (jaundice)
- Arrhythmias due to domperidone

Drug Interactions:

- Avoid concurrent use of other medications metabolized by CYP3A4 enzymes with domperidone.
- Alcohol or other CNS depressants may enhance drowsiness caused by this combination.
- Chronic use of other hepatotoxic drugs with paracetamol may increase the risk of liver damage.

Storage:

Store in a cool, dry place below 30°C. Protect from moisture and direct sunlight. Keep out of reach of children.

Packaging:

[Details of blister pack or container, depending on the manufacturer.]

Disclaimer:

This medication should only be taken under medical supervision. Consult your healthcare provider for proper diagnosis and guidance.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for