

## Paroxyle 12.5 CR

### Paroxetine 12.5mg Prolonged Release Tablets

#### An Antidepressant for Treating Mood and Anxiety Disorders

##### Composition (Per Tablet):

- **Paroxetine I.P.** – 12.5mg (Prolonged Release)

##### Product Overview:

Paroxetine is a **selective serotonin reuptake inhibitor (SSRI)** used for the treatment of **depression, anxiety disorders, obsessive-compulsive disorder (OCD)**, and other mood-related conditions. The **prolonged-release formulation** ensures a **gradual release** of the active ingredient over time, leading to **more stable blood levels** and reducing the likelihood of side effects. Paroxetine works by **increasing serotonin levels** in the brain, helping to improve mood, decrease anxiety, and manage obsessive-compulsive symptoms.

This formulation provides the **benefit of once-daily dosing**, ensuring better **patient compliance** and convenience.

##### Key Benefits:

- **Effective in treating major depressive disorder (MDD)**
- **Helps alleviate symptoms of generalized anxiety disorder (GAD) and social anxiety disorder (SAD)**
- **Reduces symptoms of obsessive-compulsive disorder (OCD)**
- **Prolonged-release** for stable blood levels and fewer side effects
- **Once-daily dosing** for ease of use
- **Well-tolerated** with a lower risk of adverse effects

##### How It Works:

Paroxetine functions by **inhibiting the reuptake of serotonin** in the brain, which increases the serotonin levels. This helps to regulate mood, improve feelings of well-being, and alleviate anxiety. The **prolonged-release formulation** helps maintain consistent therapeutic levels of the drug throughout the day, leading to **fewer peaks and troughs** in drug levels and enhancing the effectiveness and tolerability of the medication.

##### Indications:

- **Major depressive disorder (MDD)**
- **Generalized anxiety disorder (GAD)**
- **Social anxiety disorder (SAD)**
- **Obsessive-compulsive disorder (OCD)**
- **Panic disorder**
- **Post-traumatic stress disorder (PTSD)**
- **Premenstrual dysphoric disorder (PMDD)**

##### Dosage & Administration:

- **Usual Adult Dose:** 12.5mg once daily, typically taken in the morning
- The dose may be **gradually increased** based on individual response, up to 25mg daily
- **Swallow whole** with water; do not crush, chew, or break the tablet
- Can be taken **with or without food**
- **Follow your doctor's instructions** for appropriate dosage and adjustments

##### Precautions & Warnings:

- Not recommended for use in patients with **severe liver or kidney impairment**
- **Gradual dose adjustments** are advised to reduce withdrawal symptoms if discontinuing the medication
- **Monitor for signs of suicidal thoughts**, especially in children, adolescents, and young adults
- May cause **drowsiness, dizziness, or difficulty concentrating**; avoid driving or operating heavy

machinery until you know how this medication affects you

- **Serotonin syndrome** (a potentially serious condition) can occur, particularly when combined with other serotonergic drugs
- Caution in patients with **glaucoma, seizures, or bipolar disorder**
- Not recommended for use in **pregnancy** or **breastfeeding** unless deemed necessary by your healthcare provider

Manufactured in India for:



**Cafoli Lifecare Pvt. Ltd.**

**(An ISO 9001: 2015 Certified Co.)**

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for

### Side Effects (Common and Rare):

- Nausea, dry mouth, or loss of appetite
- Drowsiness, dizziness, or insomnia
- Headache or sweating
- Sexual dysfunction or changes in libido
- Weight loss or gain
- **Serotonin syndrome** (rare, serious)
- **Suicidal thoughts** (especially in younger individuals)

### Storage Instructions:

- Store in a **cool, dry place**, away from direct sunlight
- Keep out of reach of children
- Do not refrigerate or freeze

---

## Why Choose This Product?

- **Clinically proven efficacy** in treating depression, anxiety, and OCD
- **Prolonged-release formulation** for consistent, steady relief
- **Convenient once-daily dosing** for improved patient adherence
- **Better tolerability** with fewer side effects compared to immediate-release formulations

For optimal results, **follow the prescribed dosage** and regularly monitor for any side effects. If any unusual symptoms occur, seek immediate medical help.