

Information For the User
Pentoxifylline (400mg) Tablet

Pexoline PR 400 Tablet

Pexoline PR 400 Tablet

Dosage Form: Tablet

Composition:

Each film coated tablet contains:
Pentoxifylline I.P. 400 mg

Description:

Pexoline PR 400 Tablet contains Pentoxifylline, a hemorheologic agent that improves blood flow by reducing blood viscosity and enhancing microcirculation. It is indicated for the management of peripheral vascular disorders and related circulatory insufficiency. The formulation helps improve tissue oxygenation, reduce claudication symptoms, and support overall vascular health.

Indications:

Peripheral arterial disease
Intermittent claudication
Chronic occlusive arterial disease
Cerebrovascular insufficiency

Dosage & Administration:

Dosage should be taken as directed by the physician.
Tablet should be swallowed whole with water, preferably after meals.
Regular monitoring of blood pressure and renal function is recommended.

Pharmacological Action:

Pentoxifylline improves erythrocyte flexibility, decreases blood viscosity, and inhibits platelet aggregation. These effects enhance microcirculatory blood flow, oxygen delivery to tissues, and reduce symptoms associated with vascular insufficiency.

Side Effects:

Nausea
Vomiting
Headache
Dizziness
Mild gastrointestinal discomfort

Precautions & Warnings:

Use under medical supervision only.
Use cautiously in patients with hypotension, bleeding disorders, or severe cardiac conditions.
Not recommended during pregnancy and lactation unless prescribed.

Drug Interactions:

May interact with anticoagulants or antiplatelet drugs.
Consult physician before concomitant use with other medications.

Storage:

Store below 25°C in a cool, dry place.
Protect from light and moisture.
Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for