Information For the User Pregabalin I.P. 50mg + Duloxetine 20mg Capsule

Pregahike D 50/20 Capsule

Pregahike D 50/20 Capsule

Composition:

Each hard gelatin capsule contains:

- Pregabalin IP ... 50 mg
- Duloxetine Hydrochloride IP eq. to Duloxetine ... 20 mg
- Excipients ... q.s.

Dosage Form:

Capsule

Therapeutic Category:

Anticonvulsant + SNRI Antidepressant (for Neuropathic Pain & Anxiety)

Pharmacology & Mechanism of Action:

Pregabalin reduces excitatory neurotransmitter release by binding to voltage-gated calcium channels in the CNS. Duloxetine inhibits the reuptake of serotonin and norepinephrine, enhancing pain modulation and mood control. Together, they offer synergistic relief in neuropathic pain and anxiety.

Indications:

- Neuropathic pain (diabetic, sciatica, fibromyalgia)
- Generalized anxiety disorder
- Adjunctive treatment in chronic pain syndromes

Dosage & Administration:

One capsule once or twice daily or as directed by the physician Can be taken with or without food Swallow whole, do not chew

Contraindications:

- Hypersensitivity to ingredients
- Use with MAO inhibitors
- Severe hepatic or renal dysfunction

Warnings & Precautions:

- Monitor for mood or behavioral changes
- May cause drowsiness—avoid driving or machinery

- Taper dose when discontinuing
- Not advised in pregnancy/lactation unless prescribed

Adverse Effects:

- Dizziness, somnolence, nausea
- Dry mouth, constipation
- Increased BP (duloxetine), edema (pregabalin)

Drug Interactions:

- Avoid with serotonergic agents and MAOIs
- Alcohol may worsen drowsiness or liver risk
- Caution with CYP1A2/CYP2D6 inhibitors

Overdose:

Supportive care recommended. Symptoms may include agitation, confusion, or serotonin syndrome

Storage:

Store below 25°C in a cool, dry place Protect from light and moisture Keep out of reach of children

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)
Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113 TM: Trademark Applied for