

Information For the Use
Ranitidine (75mg/5ml)

Ranilair Suspension

Ranitidine Suspension (75mg/5ml)

Active Ingredient: Ranitidine Hydrochloride

Strength: 75mg per 5ml

Uses:

Ranitidine is an H₂-receptor antagonist that reduces the production of stomach acid. It is commonly used for:

- Treating and preventing heartburn.
- Relieving symptoms of gastroesophageal reflux disease (GERD).
- Treating stomach and duodenal ulcers.
- Conditions causing excessive stomach acid, such as Zollinger-Ellison syndrome.

Dosage:

- **Adults and children over 12 years:** Typically, 5–10ml (75–150mg) twice daily, or as prescribed by your doctor.
- **Children (3–11 years, depending on weight):** Doses are calculated based on body weight. Follow your doctor's guidance.

Note: Shake the bottle well before each use. Use the measuring device provided to ensure accurate dosing.

Side Effects:

Common side effects may include:

- Headache
- Constipation or diarrhea
- Nausea
- Dizziness

Rare but serious side effects:

- Allergic reactions (rash, swelling, or breathing difficulty)
- Yellowing of the skin or eyes (signs of liver issues)

Precautions:

- Do not use if you are allergic to ranitidine or any of its ingredients.
- Inform your doctor if you have kidney or liver disease.
- Not recommended for long-term use unless directed by your doctor.
- Avoid alcohol, as it may increase stomach acid and worsen symptoms.

Storage:

- Store at room temperature (20°C–25°C), away from direct light and moisture.
- Keep out of reach of children.

Warnings:

The use of ranitidine has been limited or discontinued in many regions due to concerns over impurities (NDMA). Consult your healthcare provider for alternatives or further guidance.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for