

Information For the User
Rabeprazole 40mg (EC) + Domperidone 30mg (SR)

Ribziff DSR 40 capsule

Brand Name: Ribziff DSR 40 Capsule

Dosage Form: Capsule (Enteric-Coated + Sustained Release)

Therapeutic Category: Proton Pump Inhibitor + Prokinetic Agent

Composition:

Each hard gelatin capsule contains:

- **Rabeprazole Sodium IP** (as enteric-coated pellets) equivalent to Rabeprazole – 40 mg
- **Domperidone IP** (as sustained release pellets) – 30 mg
- Excipients – q.s.
- Colours: Approved colours used in capsule shells

Pharmacology & Mechanism of Action:

Rabeprazole is a proton pump inhibitor that suppresses gastric acid secretion by inhibiting the H^+/K^+ ATPase enzyme system at the secretory surface of gastric parietal cells. It provides rapid and sustained acid suppression.

Domperidone is a dopamine D2 receptor antagonist with prokinetic and antiemetic properties. It enhances gastrointestinal motility and accelerates gastric emptying without affecting gastric secretions.

Indications:

- Gastroesophageal reflux disease (GERD)
- Non-ulcer dyspepsia
- Peptic ulcer disease
- Gastritis, epigastric pain, or bloating
- Functional dyspepsia associated with delayed gastric emptying
- Nausea and vomiting due to acid reflux or dysmotility

Dosage & Administration:

- **Adults:** One capsule once daily, preferably in the morning before food
- Dosage to be adjusted as per physician's direction
- Swallow whole – do not crush or chew

Contraindications:

- Known hypersensitivity to Rabeprazole, Domperidone, or any excipients
- Patients with prolactinoma or gastrointestinal hemorrhage, perforation, or obstruction
- Domperidone is contraindicated in patients with QT prolongation or concurrent use of QT-prolonging drugs

Warnings & Precautions:

- Use with caution in patients with hepatic or renal impairment
- Long-term use of PPIs may lead to Vitamin B12 deficiency, hypomagnesemia, or increased fracture risk
- Domperidone should be used at the lowest effective dose for the shortest duration
- Not recommended in patients with significant cardiac arrhythmias

Drug Interactions:

- Rabeprazole may reduce absorption of drugs dependent on gastric pH (e.g., ketoconazole, atazanavir)
- Domperidone's plasma levels may increase with CYP3A4 inhibitors (e.g., ketoconazole, erythromycin)
- Avoid co-administration with anticholinergics or QT-prolonging agents

Adverse Effects:

- Common: Headache, nausea, flatulence, abdominal discomfort
- Less common: Rash, dizziness, dry mouth, diarrhea
- Rare: QT prolongation (with Domperidone in high doses), gynecomastia, galactorrhea

Storage:

- Store below 25°C in a dry place
- Protect from moisture and direct light
- Keep out of reach of children

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for