

Rosumore Gold

Product Name: Rosumore Gold

Composition:

- **Aspirin I.P.** 75 mg
 - **Rosuvastatin I.P.** 10 mg
 - **Clopidogrel I.P.** 75 mg
- Dosage Form:** Tablet

Description:

Rosumore Gold is a combination tablet containing Aspirin, Rosuvastatin, and Clopidogrel. Aspirin acts as an antiplatelet agent to prevent blood clotting, while Clopidogrel further enhances platelet inhibition. Rosuvastatin works to reduce cholesterol levels, particularly low-density lipoprotein (LDL) cholesterol, in the blood. This combination therapy is used in the prevention of cardiovascular events like heart attacks and strokes.

Composition (per tablet):

- **Aspirin I.P.** – 75 mg
- **Rosuvastatin I.P.** – 10 mg
- **Clopidogrel I.P.** – 75 mg
- **Excipients** – q.s.

Indications:

Rosumore Gold is indicated for the:

1. Primary and secondary prevention of cardiovascular events (heart attack, stroke, etc.)
2. Management of atherosclerotic cardiovascular disease
3. Reducing the risk of myocardial infarction and stroke in patients with established cardiovascular diseases or at high risk
4. Prevention of blood clots post-angioplasty or stent placement

Mechanism of Action:

- **Aspirin:** Inhibits cyclooxygenase (COX) enzymes, which decreases the production of thromboxane A₂, a substance that promotes platelet aggregation.
- **Rosuvastatin:** Inhibits HMG-CoA reductase, which is involved in cholesterol synthesis, lowering LDL cholesterol and improving overall lipid profile.
- **Clopidogrel:** Inhibits ADP-induced platelet aggregation by binding to the P2Y₁₂ receptor on platelets, reducing platelet activation and clot formation.

Dosage and Administration:

- **Adult Dose:** One tablet once daily, preferably with food.
- **Special Population:** Adjust dosage for patients with hepatic or renal impairments based on the physician's recommendation.

- **Administration:** Swallow the tablet whole with a glass of water. Do not chew or crush the tablet.

Side Effects:

- **Common:** Headache, gastrointestinal discomfort, indigestion, bruising, or bleeding
- **Less Common:** Muscle pain or weakness (statin-related), liver enzyme abnormalities, allergic reactions
- **Serious:** Unusual bleeding, severe allergic reactions, or unexplained muscle pain and tenderness

Precautions:

1. **Bleeding Disorders:** Use with caution in patients with a history of bleeding disorders, peptic ulcer disease, or recent surgeries.
2. **Hepatic Impairment:** Monitor liver function tests, as Rosuvastatin can affect liver enzymes.
3. **Renal Impairment:** Dose adjustment may be necessary in patients with renal impairment.
4. **Pregnancy and Lactation:** Not recommended for use during pregnancy and breastfeeding.
5. **Gastrointestinal Issues:** May cause irritation, ulcers, or bleeding in the stomach, especially when taken on an empty stomach.

Contraindications:

- Known hypersensitivity to any of the ingredients in the formulation
- Active bleeding disorders or conditions like peptic ulcer disease
- Severe liver or renal impairment
- Pregnancy and breastfeeding

Storage:

- Store below 30°C in a cool, dry place
- Keep the container tightly closed and out of reach of children
- Protect from moisture and direct sunlight

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for