

Information For the User
Aspirin 75mg + Rosuvastatin 20mg + Clopidogrel 150mg Tablet

Rosumore Gold 20/150 Tablet

Brand Name: Rosumore Gold 20/150 Tablet

Composition:

Each film-coated tablet contains:

Rosuvastatin Calcium IP equivalent to Rosuvastatin 20 mg

Clopidogrel Bisulfate IP equivalent to Clopidogrel 150 mg

Aspirin IP 75 mg (as Gastro-resistant)

Therapeutic Indication:

Indicated for secondary prevention of cardiovascular events in patients with established atherosclerotic disease including myocardial infarction stroke and peripheral artery disease
Used in patients with acute coronary syndrome post-angioplasty or stenting and in high-risk individuals for combined lipid-lowering and antiplatelet therapy

Mechanism of Action:

Rosuvastatin is an HMG-CoA reductase inhibitor that reduces cholesterol synthesis in the liver and increases LDL clearance from plasma

Clopidogrel is a thienopyridine antiplatelet agent that irreversibly inhibits the P2Y₁₂ receptor on platelets preventing aggregation

Aspirin is an antiplatelet agent that irreversibly inhibits cyclooxygenase-1 (COX-1) and thromboxane A₂ production reducing platelet aggregation

The combination works synergistically to prevent thrombus formation and manage dyslipidemia thereby reducing the risk of recurrent cardiovascular events

Dosage & Administration:

Usual adult dose: one tablet once daily or as directed by the physician

Should be taken preferably after meals to minimize gastric irritation

Swallow whole with water do not crush or chew

Contraindications:

Hypersensitivity to any component of the formulation

Active bleeding disorders such as peptic ulcer or intracranial hemorrhage

Severe hepatic or renal impairment

History of aspirin-induced asthma or NSAID allergy

Pregnancy and lactation

Warnings & Precautions:

Monitor lipid profile liver function platelet count and bleeding parameters periodically

Use with caution in patients with history of gastrointestinal bleeding or clotting disorders

Avoid alcohol and concurrent use of other NSAIDs

Discontinue at least 5–7 days prior to surgery if antiplatelet effect is not desired

Side Effects:

Common: headache myalgia gastritis dyspepsia dizziness

Less common: rash elevated liver enzymes minor bleeding bruising

Rare: gastrointestinal bleeding rhabdomyolysis thrombocytopenia hypersensitivity reactions

Storage:

Store in a cool dry and dark place below 25°C

Protect from moisture and direct sunlight

Keep out of reach of children

Packaging:

Rosumore Gold 20/150 Tablet is available in alu-alu strip packs

Schedule:

Schedule H Prescription Drug – To be sold by retail on the prescription of a registered medical practitioner only

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for