#### Information For the User Sacubitril 49 mg + Valsartan 51 mg tablet

# Sacustep 100 Tablet

#### **Composition:**

Each film-coated bilayered tablet contains: Sacubitril – 49 mg Valsartan – 51 mg Excipients – q.s.

## **Description:**

Sacustep 100 Tablet is a fixed-dose combination of Sacubitril and Valsartan, classified as an angiotensin receptor neprilysin inhibitor (ARNI). This formulation is indicated for the treatment of chronic heart failure with reduced ejection fraction (HFrEF), offering improved survival and reduced hospitalization compared to traditional therapies. Sacubitril enhances the levels of beneficial natriuretic peptides, while Valsartan counteracts the harmful effects of the renin-angiotensin-aldosterone system (RAAS).

## **Indications:**

Chronic heart failure (NYHA Class II–IV) with reduced ejection fraction Alternative to ACE inhibitors or ARBs in stable symptomatic heart failure patients Used to reduce cardiovascular mortality and morbidity

## **Mechanism of Action:**

Sacubitril is a neprilysin inhibitor that prevents the degradation of natriuretic peptides, bradykinin, and other vasoactive peptides. This leads to vasodilation, natriuresis, diuresis, and inhibition of cardiac remodeling.

Valsartan blocks the angiotensin II type 1 (AT1) receptor, reducing vasoconstriction, sodium retention, and aldosterone secretion. Together, the combination improves hemodynamics and delays progression of heart failure.

# **Dosage and Administration:**

**Recommended dose:** One tablet twice daily (Sacubitril 49 mg + Valsartan 51 mg) Patients should be stabilized on lower doses before up-titration to Sacustep 100 Administer orally, with or without food Ensure a 36-hour washout period after discontinuing ACE inhibitors before initiating therapy

# Side Effects:

Dizziness, hypotension Hyperkalemia Renal function impairment Cough Rare: Angioedema

## **Precautions:**

Not to be used with ACE inhibitors or in patients with a history of angioedema Monitor blood pressure, renal function, and potassium levels regularly Contraindicated during pregnancy and lactation Use with caution in patients with hepatic or renal impairment

#### **Storage:**

Store below 25°C in a dry place Protect from direct sunlight and moisture Keep out of reach of children

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd. (An ISO 9001: 2015 Certified Co.) Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113 TM: Trademark Applied for