Dicyclomine HCL 10mg + Diclofenac Sodium 25mg

Scotspas D Injection

Dicyclomine HCL (10 mg) + Diclofenac Sodium (25 mg) Injection

Description:

Dicyclomine HCL (10 mg) + Diclofenac Sodium (25 mg) injection is a combination of Dicyclomine, an antispasmodic, and Diclofenac Sodium, a nonsteroidal anti-inflammatory drug (NSAID). This combination is used to provide relief from acute pain and discomfort associated with gastrointestinal and musculoskeletal conditions.

Composition:

• Dicyclomine HCL: 10 mg

• Diclofenac Sodium: 25 mg

Indications:

This combination injection is indicated for the following conditions:

- 1. Irritable Bowel Syndrome (IBS):
 - For the relief of abdominal cramps and pain caused by IBS.
- 2. Abdominal Pain and Discomfort:
 - For treating pain and discomfort related to gastrointestinal spasms, such as those caused by colitis or functional dyspepsia.
- 3. Postoperative Pain:
 - For pain management following abdominal or gastrointestinal surgery, especially where muscle spasms and inflammation are involved.

Mechanism of Action:

• Dicyclomine (10 mg):
Dicyclomine is an anticholinergic agent that works by blocking the action of

acetylcholine on smooth muscles in the gastrointestinal tract. Diclofenac Sodium

(25 mg):

Diclofenac is a NSAID that works by inhibiting cyclooxygenase (COX) enzymes, particularly COX-2, which are involved in the production of prostaglandins—chemicals in the body that cause inflammation, pain, and fever.

Dosage and Administration:

- Adults:
 - The typical dosage for Dicyclomine 10 mg + Diclofenac Sodium 25 mg injection is administered by deep intramuscular injection, usually as a one-time dose or as directed by a healthcare provider.
 - For abdominal cramps and spasms: One injection may be sufficient for relief, but further doses may be given depending on clinical response.
- Children:
 - This combination injection is typically not recommended for use in children due to the potential risk of side effects, especially for Diclofenac.

Note:

 The injection should be administered by a healthcare professional in a clinical setting. It should not be selfadministered.

Side Effects:

• Common:

 Dicyclomine: Dry mouth, dizziness, blurred vision, constipation, and urinary retention.

Serious but Rare:

- Cardiovascular issues: High blood pressure, heart attack, or stroke (especially with long-term use of Diclofenac).
- Gastrointestinal problems: Risk of ulcers, bleeding, or perforation in the stomach or intestines, particularly in elderly patients or with long-term use of Diclofenac.

Precautions:

1. Cardiovascular Conditions:

 Use with caution in patients with a history of heart disease, high blood pressure, or stroke, as Diclofenac can increase the risk of cardiovascular events, particularly with long-term use.

2. Gastrointestinal Disorders:

 Use with caution in patients with a history of ulcers, gastritis, or GI bleeding, as Diclofenac can exacerbate these conditions.

3. Liver and Kidney Impairment:

 Both Dicyclomine and Diclofenac should be used with caution in patients with liver or kidney disease. Regular monitoring of liver and kidney function may be required, particularly with prolonged treatment.

4. Pregnancy and Breastfeeding:

 Pregnancy: This medication should only be used during pregnancy if absolutely necessary and prescribed by a doctor, particularly in the third trimester, due to potential risks to the fetus.

5. Elderly Patients:

Older adults may be more susceptible to the side effects of both Dicyclomine (e.g., confusion, dry mouth) and Diclofenac (e.g., gastrointestinal bleeding), requiring careful monitoring.

Storage:

- Store at room temperature (15-30°C), away from direct sunlight and moisture.
- Do not freeze.
- Keep the vial tightly sealed and out of reach of children.
- Discard any unused solution properly according to local disposal guidelines.

Note:

This combination injection should be administered only under the supervision of a healthcare provider.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.) Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for