

# Scotspas P Tablet

**Brand Name:** Scotspas P Tablet

**Dosage Form:** Tablet

**Therapeutic Category:** Antispasmodic with Analgesic

## Composition:

Each uncoated tablet contains:

- **Dicyclomine Hydrochloride IP** – 20 mg
- **Paracetamol IP** – 325 mg
- Excipients – q.s.

## Description:

Scotspas P Tablet is a synergistic combination of an anticholinergic agent and an analgesic-antipyretic. Dicyclomine provides antispasmodic action by relaxing smooth muscles of the gastrointestinal tract, while Paracetamol relieves associated pain and fever. This dual-action makes Scotspas P effective in managing spasmodic abdominal pain of gastrointestinal, biliary, or genitourinary origin.

## Pharmacology & Mechanism of Action:

- **Dicyclomine:** Acts as an anticholinergic by blocking muscarinic receptors on smooth muscle, leading to muscle relaxation and relief from spasms.
- **Paracetamol:** Inhibits prostaglandin synthesis in the CNS, providing analgesic and antipyretic effects.

## Indications:

- Abdominal colic and cramps
- Gastrointestinal spasm
- Irritable bowel syndrome (IBS)
- Dysmenorrhea
- Spastic conditions of the urinary or biliary tract

## Dosage & Administration:

- Adults: 1 tablet 2–3 times daily after meals or as directed by the physician
- Not recommended in children unless prescribed
- Do not exceed the prescribed dosage

## Contraindications:

- Hypersensitivity to dicyclomine, paracetamol, or any excipients
- Glaucoma, myasthenia gravis, obstructive uropathy, or gastrointestinal obstruction
- Severe hepatic impairment

## Warnings & Precautions:

- Use with caution in patients with hepatic or renal impairment
- Avoid alcohol and other hepatotoxic agents during treatment
- Prolonged use should be monitored to avoid hepatotoxicity from paracetamol
- Not recommended during pregnancy and lactation unless advised by a physician

**Drug Interactions:**

- Alcohol and other CNS depressants may potentiate dicyclomine's anticholinergic effects
- Increased risk of liver toxicity with excessive paracetamol or in combination with enzyme inducers

**Adverse Effects:**

- Drowsiness, dry mouth, dizziness, constipation
- In rare cases: hepatotoxicity (due to paracetamol), blurred vision, or urinary retention

**Storage:**

- Store below 25°C in a dry, dark place
- Protect from light and moisture
- Keep out of reach of children

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,  
Panchkula-134113

TM: Trademark Applied for