

Information For the User
Amisulpride (50mg) Tablet

Sulpride 50 Tablet

Brand Name: Sulpride 50 Tablet

Dosage Form: Tablet

Therapeutic Category: Antipsychotic – Atypical/Selective Dopamine Antagonist

Composition:

Each tablet contains:

- Amisulpride I.P. 50 mg
- Excipients q.s.

Pharmacology & Mechanism of Action:

Amisulpride is a substituted benzamide antipsychotic that selectively antagonizes dopamine D2 and D3 receptors in the central nervous system, particularly in the limbic system. At lower doses, it provides antipsychotic and antidepressant effects with minimal extrapyramidal side effects. By modulating dopaminergic transmission, it helps alleviate symptoms of schizophrenia and other psychotic disorders, including hallucinations, delusions, and thought disturbances.

Indications:

- Schizophrenia – mild to moderate cases
- Psychotic disorders with predominant positive symptoms
- Adjunct therapy for mood disorders with psychotic features

Dosage & Administration:

- Adults: 50–200 mg per day, divided into 1–2 doses, or as prescribed by the physician
- Tablets should be taken orally, with or without food
- Dose adjustments may be required for elderly patients or those with renal/hepatic impairment

Contraindications:

- Hypersensitivity to Amisulpride or any tablet components
- Severe renal or hepatic impairment
- History of prolactin-dependent tumors

Warnings & Precautions:

- Monitor for extrapyramidal symptoms and tardive dyskinesia
- Use cautiously in patients with cardiovascular disorders, seizures, or hyperprolactinemia
- Not recommended for children under 18 without specialist supervision
- Avoid abrupt withdrawal; taper gradually under medical guidance

Adverse Effects:

- Very common: Insomnia, anxiety, agitation
- Common: Dizziness, headache, weight gain, gastrointestinal upset
- Rare: Extrapyramidal symptoms, hyperprolactinemia, galactorrhea, gynecomastia

Drug Interactions:

- CNS depressants may enhance sedation
- Concomitant dopamine antagonists should be used cautiously under medical supervision

Overdose:

- Symptoms: Severe sedation, hypotension, extrapyramidal reactions, cardiac arrhythmias
- Treatment: Supportive care; gastric lavage if ingestion is recent

Storage:

Store below 25°C in a cool, dry place.

Protect from light and moisture.

Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for