

Tadafly 10 Tablet

Tadafly 10 Tablet: Comprehensive Information

Generic Name: Tadalafil

Brand Name: Tadafly 10

Manufacturer: [Insert Manufacturer Information, if available]

Strength: 10mg

Tadafly 10 Tablet contains Tadalafil as its active ingredient. It is widely used to treat:

- Erectile Dysfunction (ED):** Helps men achieve and maintain an erection during sexual activity.
- Benign Prostatic Hyperplasia (BPH):** Relieves symptoms like difficulty urinating or frequent urination.
- Combination of ED and BPH:** Addresses both conditions simultaneously.

Key Features of Tadafly 10

- Class of Drug:** Phosphodiesterase-5 (PDE5) Inhibitor.
- Mechanism of Action:** Inhibits PDE5, enhancing blood flow by relaxing smooth muscles in blood vessels.

Indications and Uses

- Erectile Dysfunction (ED):**
 - Helps improve erectile function when sexually stimulated.
- Benign Prostatic Hyperplasia (BPH):**
 - Relieves symptoms such as difficulty starting urination, weak stream, or frequent urge to urinate, especially at night.
- Pulmonary Arterial Hypertension (PAH):**
 - At higher doses (not 10mg), improves exercise capacity.

Dosage and Administration

- For Erectile Dysfunction:**
 - On-Demand Use:** One Tadafly 10 tablet taken at least 30 minutes before sexual activity.
 - Maximum Frequency:** Once per day.
- For Benign Prostatic Hyperplasia:**
 - Not typically prescribed at 10mg; a lower daily dose (5mg) is more common.
- For Combined ED and BPH:**
 - Dose and regimen determined by the healthcare provider.
- Administration:**
 - Take orally with water.
 - Can be taken with or without food.

Precautions Before Taking Tadafly 10

Do not take Tadafly 10 if you:

- Use nitrates (e.g., nitroglycerin) for heart conditions.
- Have a history of severe allergic reactions to Tadalafil or other PDE5 inhibitors.
- Have severe heart disease, liver impairment, or very low blood pressure.

Inform your doctor if you have:

- Cardiovascular conditions (e.g., recent heart attack or stroke).
- Vision problems, such as retinitis pigmentosa or NAION.
- Renal impairment.
- A deformed penis shape (Peyronie's disease).
- A history of priapism (prolonged erections).

Potential Side Effects

- Common Side Effects:**

- Headache, flushing, nasal congestion, indigestion, back pain, or muscle aches.
- 2. **Uncommon or Serious Side Effects:**
 - Sudden loss of vision or hearing.
 - Chest pain, irregular heartbeat, or fainting.
- 3. **Rare Side Effects:**
 - Severe allergic reactions (rash, itching, swelling).
 - Symptoms of a heart attack (chest pain, nausea, sweating).

- Keep away from moisture, heat, and direct sunlight.
- Keep out of reach of children.

Note: Always use Tadafly 10 under the guidance of a healthcare professional. Discuss your full medical history and all medications you are taking to ensure safety and efficacy.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for

Drug Interactions

Tadafly 10 may interact with:

- **Nitrates:** Can cause severe hypotension.
- **Alpha-Blockers:** Risk of low blood pressure.
- **Antifungals and Antibiotics:** Such as ketoconazole, erythromycin, or ritonavir.

Substances to Avoid:

- Grapefruit juice: May increase Tadalafil levels, heightening side effects.
- Excessive alcohol: Can enhance side effects like dizziness and low blood pressure.

Overdose Management

Taking more than the prescribed dose may result in severe side effects such as:

- Prolonged erections.
- Severe headaches or dizziness.
- Fainting or chest pain.

In case of overdose, seek immediate medical attention.

Storage Guidelines

- Store at room temperature (20–25°C / 68–77°F).