

Diziderm MS Cream

Diziderm MS Cream
(Metronidazole 1% w/w + Sucralfate 7% w/w + Povidone Iodine 5% w/w Ointment)

Description:

Diziderm MS Cream is a topical formulation designed for effective wound care and treatment of infected skin conditions. It combines the antibacterial, antifungal, and antiseptic properties of Metronidazole, Sucralfate, and Povidone Iodine to promote healing, prevent infection, and protect the affected area.

Composition:

Each 1 gram of cream contains:

- **Metronidazole (1% w/w):** An antimicrobial agent that fights anaerobic bacteria and protozoa, reducing infection and inflammation.
- **Sucralfate (7% w/w):** A cytoprotective agent that forms a protective barrier over wounds, promoting tissue regeneration and healing.
- **Povidone Iodine (5% w/w):** A broad-spectrum antiseptic that kills bacteria, fungi, and viruses, preventing and controlling infections.

Indications:

- Management of infected wounds and ulcers.
- Treatment of surgical wounds, pressure sores, and burns.
- Prevention and control of secondary infections in dermal conditions.

Mechanism of Action:

1. **Metronidazole:** Disrupts bacterial DNA synthesis, effectively killing anaerobic bacteria and protozoa.
2. **Sucralfate:** Forms a viscous protective layer on the wound surface, shielding it from external contaminants and aiding in tissue repair.

3. **Povidone Iodine:** Releases iodine, which rapidly kills pathogens on contact, ensuring a sterile wound environment.

Directions for Use:

- Clean and dry the affected area thoroughly.
- Apply a thin layer of Diziderm MS Cream to the area 2–3 times daily or as directed by a healthcare professional.
- Cover with sterile gauze if required.

Precautions:

- For external use only. Avoid contact with eyes, nose, and mouth.
- Discontinue use if irritation, redness, or swelling persists or worsens.
- Consult a healthcare professional before use during pregnancy or breastfeeding.

Contraindications:

- Hypersensitivity to Metronidazole, Sucralfate, Povidone Iodine, or any excipient in the formulation.
- Not recommended for use on large wounds or deep puncture wounds without medical advice.

Side Effects:

- Mild irritation, redness, or itching at the application site.
- Rarely, allergic reactions such as rash or swelling may occur.

Storage:

- Store in a cool, dry place below 25°C.
- Keep out of reach of children.

Presentation:

Available in a convenient ointment tube for easy application.

Caution:

**Use only as directed by a healthcare professional.
If symptoms persist, consult your doctor.**

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for