

Information For the Use of
Travoprost (0.004% w/v) Eye Drops

Travopine Eye Drops

Brand Name: Travopine Eye Drops

Dosage Form: Eye Drops

Composition:

Each ml contains:

- Travoprost IP 0.004% w/v
- Sterile ophthalmic vehicle q.s.

Description:

Travopine Eye Drops contains Travoprost, a prostaglandin F_{2α} analogue indicated for the reduction of elevated intraocular pressure in patients with open-angle glaucoma or ocular hypertension. It helps in lowering intraocular pressure by increasing the outflow of aqueous humor from the eye, thereby reducing the risk of optic nerve damage and progressive vision loss associated with glaucoma.

Indications:

Indicated for the treatment of:

- Open-angle glaucoma
- Ocular hypertension
- Elevated intraocular pressure requiring prostaglandin analogue therapy

It helps in reducing intraocular pressure and preserving visual function by minimizing glaucoma-related optic nerve damage.

Dosage & Administration:

Instill one drop into the affected eye(s) once daily in the evening or as directed by the physician.

Do not exceed the recommended dosing frequency, as this may reduce the pressure-lowering effect.

Avoid touching the dropper tip to any surface to prevent contamination.

If using multiple ophthalmic preparations, maintain an interval of at least 5–10 minutes between applications.

Pharmacological Action:

- Travoprost is a prostaglandin F_{2α} analogue that increases uveoscleral outflow of aqueous humor from the eye.
- It effectively lowers intraocular pressure without significantly affecting aqueous humor production.

These actions help in sustained reduction of elevated intraocular pressure and protection of optic nerve function.

Side Effects:

Common side effects include conjunctival hyperemia, eye irritation, itching, foreign body sensation, and blurred vision.

Long-term use may cause gradual darkening of the iris, increased eyelash growth, or pigmentation of the eyelid skin.

Rarely, ocular inflammation may occur.

Precautions & Warnings:

- For ophthalmic use only
- Remove contact lenses before administration and reinsert after at least 15 minutes
- Use with caution in patients with a history of uveitis or macular edema
- Inform patients about the possibility of permanent iris pigmentation changes
- Avoid contamination of the dropper tip

Drug Interactions:

No significant systemic drug interactions are expected due to minimal systemic absorption.

May be used concomitantly with other ophthalmic antiglaucoma medications as directed by the physician.

Storage:

Store below 25°C in a cool, dry place.

Protect from light.

Keep the bottle tightly closed when not in use.

Keep out of reach of children.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I, Panchkula-134113

Corporate Office :

Plot no. 307, Industrial Area Phase-2,
Panchkula, Haryana -134113

TM: Trademark Applied for