

Traxlink ET Tablet

Information for the Use

Tranexamic Acid (250 mg) + Ethamsylate (250 mg) Tablet

Composition:

Each tablet contains:

- **Tranexamic Acid: 250 mg**
 - **Ethamsylate: 250 mg**
-

Description:

Tranexamic Acid is an antifibrinolytic agent that helps reduce bleeding by preventing the breakdown of blood clots. Ethamsylate is a hemostatic agent that helps in stabilizing blood vessels, reducing capillary bleeding, and improving platelet function.

Indications:

Tranexamic Acid (250 mg) + Ethamsylate (250 mg) Tablet is indicated for the treatment and prevention of excessive bleeding in the following conditions:

- **Menorrhagia:** Heavy menstrual bleeding.
 - **Post-surgical bleeding:** To prevent or reduce bleeding after surgery.
 - **Dental or nasal bleeding:** Particularly in patients with clotting disorders.
 - **Trauma or injury:** In cases of significant blood loss.
-

Mechanism of Action:

- **Tranexamic Acid** works by inhibiting fibrinolysis (the breakdown of blood clots), allowing the clots to remain intact and reduce bleeding.
 - **Ethamsylate** helps stabilize capillaries, improve platelet aggregation, and enhance blood clotting by strengthening blood vessels and reducing the risk of bleeding from small blood vessels.
-

Dosage and Administration:

- **Adult Dosage:**
 - The usual dose is 1 tablet (250 mg Tranexamic Acid + 250 mg Ethamsylate), 2 to 3 times daily, or as directed by the healthcare provider.
 - The dose may vary depending on the severity of the bleeding or the medical condition being treated.
- **Pediatric Dosage:**
 - The use of this combination in children should be determined by a healthcare provider based on the child's age, weight, and medical condition.

Directions for Use:

- Swallow the tablet whole with a glass of water.
 - Can be taken with or without food.
 - Follow the prescribed dosage and duration as directed by your healthcare provider.
-

Contraindications:

- **Hypersensitivity to Tranexamic Acid, Ethamsylate, or any other components of the tablet.**
 - **Active thromboembolic disease (such as deep vein thrombosis, pulmonary embolism, or stroke).**
 - **History of or active blood clotting disorders unless specifically prescribed under medical supervision.**
-

Precautions:

- **Renal Impairment: Use with caution in patients with impaired kidney function.**
 - **Pregnancy and Lactation:**
 - **Pregnancy: Should only be used during pregnancy if the benefits outweigh the risks, and after careful consideration by a healthcare provider.**
-

Possible Side Effects:

Common side effects:

- **Nausea or vomiting.**
 - **Diarrhea or constipation.**
 - **Headache or dizziness.**
 - **Mild skin rash or itching.**
-

Serious side effects (seek medical attention immediately):

- **Thromboembolic events: Signs of blood clots, such as swelling or pain in the legs, difficulty breathing, or chest pain.**
 - **Gastrointestinal bleeding: Black or tarry stools, or blood in the vomit.**
 - **Severe skin reactions: Rash, blistering, or peeling.**
-

Drug Interactions:

- **Oral contraceptives: May increase the risk of thrombosis (blood clots), use with caution.**
 - **Anticoagulants (e.g., warfarin, heparin): Use cautiously, as there may be an increased risk of bleeding or clotting.**
 - **Other hemostatic drugs: Caution is advised when used with other medications that affect blood clotting.**
-

Storage:

- **Store in a cool, dry place at 20°C to 25°C (68°F to 77°F), away from direct sunlight and moisture.**
 - **Keep out of the reach of children.**
-

Special Instructions:

- **Take the full prescribed course, even if symptoms improve before finishing the medication.**
 - **Do not exceed the prescribed dosage.**
 - **If you experience any unusual symptoms, such as chest pain, swelling, or difficulty breathing, seek medical attention immediately.**
-

Note: This medication should be used under the supervision of a healthcare provider. Always consult your doctor if you have any concerns or if you experience any side effects during treatment with Tranexamic Acid (250 mg) + Ethamsylate (250 mg) Tablet.

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for