

Traxlink MF tablet

Information for the Use

Tranexamic Acid (500 mg) + Mefenamic Acid (250 mg) Tablet

Composition:

Each tablet contains:

- **Tranexamic Acid: 500 mg**
 - **Mefenamic Acid: 250 mg**
-

Description:

Tranexamic Acid is an antifibrinolytic agent that helps prevent the breakdown of blood clots, reducing excessive bleeding. Mefenamic Acid is a non-steroidal anti-inflammatory drug (NSAID) that works by reducing inflammation, pain, and fever. The combination of Tranexamic Acid and Mefenamic Acid is used to treat conditions involving excessive bleeding while providing pain relief, particularly in cases of heavy menstrual bleeding or other bleeding disorders.

Indications:

This combination tablet is indicated for the following conditions:

- **Menorrhagia: Heavy menstrual bleeding, to reduce bleeding and control associated pain.**
- **Post-operative bleeding: To reduce bleeding after surgery.**
- **Trauma or injury: To manage bleeding and associated pain following an injury.**
- **Dental and nasal bleeding: To reduce excessive bleeding.**

- **Other bleeding disorders: Where both excessive bleeding and pain need to be managed simultaneously.**
-

Mechanism of Action:

- **Tranexamic Acid works by inhibiting the breakdown of fibrin in blood clots, thereby promoting clot stability and reducing bleeding.**
 - **Mefenamic Acid is an NSAID that inhibits cyclooxygenase (COX) enzymes, which are responsible for producing prostaglandins that mediate pain, inflammation, and fever. This helps in reducing pain and inflammation while preventing excessive bleeding.**
-

Dosage and Administration:

- **Adult Dosage:**
 - **The typical dose is 1 tablet (500 mg Tranexamic Acid + 250 mg Mefenamic Acid) taken 2 to 3 times daily for up to 5 days during menstruation, or as directed by your healthcare provider.**
 - **For post-surgical bleeding or other conditions, your doctor will provide a specific dosing regimen based on the severity of the condition.**
- **Pediatric Dosage:**
 - **Use in children should be prescribed by a healthcare provider after evaluating the child's age, weight, and medical condition.**

Directions for Use:

- **Swallow the tablet whole with a glass of water.**
 - **Can be taken with or without food.**
 - **Follow your healthcare provider's dosage instructions carefully.**
-

Contraindications:

- **Hypersensitivity to Tranexamic Acid, Mefenamic Acid, or any other components of the tablet.**
 - **Severe renal impairment or history of kidney disease.**
 - **Pregnancy (especially in the third trimester) and breastfeeding, unless advised by a healthcare provider.**
 - **Children under 12 years of age.**
-

Precautions:

- **Renal Impairment:** Use with caution in patients with kidney disease, as both Tranexamic Acid and Mefenamic Acid are excreted by the kidneys.
 - **Bleeding Disorders:** Caution should be exercised in patients with clotting disorders or a history of thromboembolic disease.
 - **Pregnancy and Lactation:**
 - **Pregnancy:** Not recommended, especially in the third trimester, due to potential risks to the fetus.
 - **Breastfeeding:** Caution is advised as both drugs may pass into breast milk. Consult your doctor before use during lactation.
-

Possible Side Effects:

Common side effects:

- **Stomach upset, nausea, or vomiting.**
- **Diarrhea or constipation.**
- **Dizziness or headache.**
- **Mild rash or itching.**

Serious side effects (seek medical attention immediately):

- **Severe gastrointestinal symptoms:** Persistent stomach pain, black or tarry stools, or vomiting blood.
 - **Thromboembolic events:** Signs of a blood clot, such as pain or swelling in the legs, chest pain, or shortness of breath.
-

Drug Interactions:

- **Anticoagulants (e.g., warfarin):** The combination of Tranexamic Acid and Mefenamic Acid may increase the risk of bleeding.
 - **Other NSAIDs:** Use with caution as it may increase the risk of gastrointestinal side effects like ulcers or bleeding.
-

Storage:

- **Store at 20°C to 25°C (68°F to 77°F), away from heat and moisture.**
 - **Keep the medication out of the reach of children.**
-

Note: This medication should be used under the supervision of a healthcare professional. Always consult your healthcare provider for any concerns or questions about the use of **Tranexamic Acid (500 mg) + Mefenamic Acid (250 mg) Tablet.**

Manufactured in India for:

CafoliTM
L I F E C A R E

Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113

TM: Trademark Applied for