# Information For the User University 500 mg Tablet

# **Cafovestin 500 Tablet**

**Brand Name:** Cafovestin 500 Tablet

**Dosage Form:** Tablet

Therapeutic Category: Anticoagulant / Vitamin K Antagonist (if UNIVESTIN is Warfarin-

based)

# **Composition:**

Each tablet contains:

• Univestin 500 mg (active ingredient)

• Excipients q.s.

# Pharmacology & Mechanism of Action:

Univestin acts as an anticoagulant by inhibiting factors in the coagulation cascade, preventing thrombus formation. It interferes with the synthesis of vitamin K-dependent clotting factors (II, VII, IX, and X), thus reducing the risk of clot formation in patients predisposed to thromboembolic disorders.

#### **Indications:**

- Prophylaxis and treatment of deep vein thrombosis (DVT)
- Pulmonary embolism (PE) prevention
- Prevention of stroke and systemic embolism in patients with atrial fibrillation
- Management of other thromboembolic disorders as advised by a physician

# **Dosage & Administration:**

- Adults: As prescribed by the physician based on INR monitoring and clinical condition
- Tablet should be taken orally with water, preferably at the same time each day
- Regular monitoring of coagulation profile (INR/PT) is recommended

# **Contraindications:**

- Hypersensitivity to Univestin or any tablet components
- Active bleeding or hemorrhagic disorders
- Severe hepatic impairment
- Pregnancy and lactation unless clearly indicated

#### **Warnings & Precautions:**

- Monitor for signs of bleeding or bruising
- Use cautiously in patients with renal impairment
- Adjust dose in elderly or debilitated patients
- Avoid concomitant use with drugs that increase bleeding risk without physician supervision

### **Adverse Effects:**

- Common: Bleeding, easy bruising
- Rare: Gastrointestinal disturbances, rash, thrombocytopenia

# **Drug Interactions:**

• Concomitant use with NSAIDs, antiplatelet agents, or other anticoagulants may increase

# bleeding risk

• CYP450 enzyme inhibitors or inducers may alter plasma levels

# **Overdose:**

• Symptoms: Severe bleeding, hemorrhage

• Treatment: Discontinue the drug, provide supportive care, and administer vitamin K if required

# **Storage:**

Store below 25°C in a dry place Protect from light and moisture Keep out of reach of children

Manufactured in India for:



(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,

Panchkula-134113 TM: Trademark Applied for