

Uriviga DFZ Capsule

Tamsulosin 0.4 mg + Deflazacort 30 mg Capsule

Dual-Action Solution for Symptomatic Relief in Benign Prostatic Hyperplasia (BPH) with Inflammatory Conditions

This combination capsule of Tamsulosin and Deflazacort is formulated to manage the symptoms of Benign Prostatic Hyperplasia (BPH) while addressing inflammation-related complications. The blend of an alpha-blocker and a corticosteroid provides relief from urinary obstruction and reduces inflammation, improving the quality of life in men with BPH and associated inflammatory conditions.

Composition & Mechanism of Action

- Tamsulosin (0.4 mg)**
 - Class:** Alpha-1 adrenergic receptor blocker.
 - Action:** Relaxes the smooth muscles of the bladder neck and prostate, improving urinary flow and relieving symptoms like frequent urination and difficulty in starting urination.
- Deflazacort (30 mg)**
 - Class:** Corticosteroid.
 - Action:** Reduces inflammation, edema, and immune responses associated with BPH or other coexisting inflammatory conditions. Its immunosuppressive properties help manage swelling and discomfort effectively.

Key Benefits

- Improves Urinary Flow:** Tamsulosin provides targeted relief from urinary obstruction and improves bladder emptying.
- Reduces Inflammation:** Deflazacort addresses inflammation-related discomfort, swelling, and other associated symptoms.

- Dual Symptom Management:** Combines relief from urinary issues with anti-inflammatory benefits, especially useful for BPH with coexisting conditions.
- Non-Invasive Relief:** Helps manage symptoms without the immediate need for surgical intervention.

Indications

- Benign Prostatic Hyperplasia (BPH) with associated inflammatory conditions.
- Inflammatory complications or edema in the genitourinary tract.

Dosage and Administration

- Recommended Dose:** One capsule daily or as prescribed by your healthcare provider.
- Take the capsule after meals to minimize gastric discomfort.
- Swallow whole with water, preferably at the same time each day.

Precautions

- Monitor Blood Pressure:** Tamsulosin may lower blood pressure; caution is advised in patients prone to hypotension.
- Corticosteroid Precautions:** Long-term use of Deflazacort can suppress the immune system and may require monitoring for infections.
- Avoid Sudden Discontinuation:** Do not stop taking Deflazacort abruptly without consulting a doctor, as it may lead to adrenal insufficiency.
- Rule Out Prostate Cancer:** Ensure prostate cancer is excluded before starting treatment, as symptoms may overlap.
- Allergic Reactions:** Discontinue use if signs of hypersensitivity occur, such as rash, swelling, or difficulty breathing.

Potential Side Effects

- **Common:** Dizziness, headache, nausea, and increased appetite.
- **Corticosteroid-Specific:** Weight gain, mood changes, fluid retention, and mild skin thinning.
- **Less Common:** Visual disturbances, muscle weakness, or severe allergic reactions—seek medical attention if these occur.

Additional Information

- **Storage:** Keep in a cool, dry place, away from direct sunlight and moisture.
- **Packaging:** Available in blister packs of 10 or 30 capsules.
- **Shelf Life:** Best before 24 months from the date of manufacture.

Experience comprehensive relief from BPH and associated inflammation with Tamsulosin + Deflazacort Capsules. Consult your healthcare provider to ensure this dual-action therapy is the right choice for you.

Manufactured in India for:



Cafoli Lifecare Pvt. Ltd.

(An ISO 9001: 2015 Certified Co.)

Plot no.: 367-FF, Industrial Area Phase-I,
Panchkula-134113

TM: Trademark Applied for